

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 June 2003 (05.06.2003)

PCT

(10) International Publication Number
WO 03/045988 A2

(51) International Patent Classification⁷: **C07K 14/39**,
C12N 15/10

(21) International Application Number: PCT/EP02/13549

(22) International Filing Date:
28 November 2002 (28.11.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
01204600.9 29 November 2001 (29.11.2001) EP

(71) Applicant (*for all designated States except US*):
**VLAAMS INTERUNIVERSITAIR INSTITUUT
VOOR BIOTECHNOLOGIE VZW** [BE/BE]; Rijkswis-
chestraat 120, B-9052 Zwijnaarde (BE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **CONTRERAS,
Roland, Henry** [BE/BE]; Molenstraat 53, B-9820 Merel-
beke (BE). **CHEN, Cuiying** [BE/BE]; Lijsterstraat 10,
B-9860 Balegem (BE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,
SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US,
UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished
upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: METHOD TO ISOLATE GENES INVOLVED IN AGING

(57) Abstract: The present invention relates to a method to isolate genes involved in aging and/or oxidative stress, by mutation or transformation of a yeast cell, subsequent screening of the mutant or transformed cells that are affected in aging and isolation of the affected gene or genes, and the use of these genes to modulate aging and aging-associated diseases in a eukaryotic cell and/or organism.

BEST AVAILABLE COPY

METHOD TO ISOLATE GENES INVOLVED IN AGING

The present invention relates to a method to isolate genes involved in aging and/or aging- associated diseases and/or oxidative stress, by mutation or transformation of a yeast cell, subsequent screening of the mutant or transformed cells that are affected in aging and isolation of the affected gene or genes, and the use of these genes to modulate aging and aging-associated diseases in a eukaryotic cell and/or organism.

Aging is a process in which all individuals of a species undergo a progressive decline in vitality leading to aging-associated diseases (AAD's) and to death. The process of aging is influenced by many factors, including metabolic capacity, stress resistance, genetic stability and gene regulation (Jazwinski, 1996). The final life span of an organism is also affected by the sum of deleterious changes and counteracting repair and maintenance mechanisms (Johnson *et al.*, 1999).

Several approaches have been followed to study aging. These include the identification of key genes and pathways important in aging, the study of genetic heritable diseases associated with aging, physiological experiment and advanced molecular biology studies of model organisms. Among these organisms, *Caenorhabditis elegans*, *Drosophila melanogaster* and the budding yeast *Saccharomyces cerevisiae* have a life span that can be influenced by single gene mutations or overexpression of a particular protein (Johnson *et al.*, 1999). Especially *S. cerevisiae* has been used as one of the model organisms to study the aging process (Gershon and Gershon, 2000). Yeast life span is defined as the number of daughter cells produced by mother cells before they stop dividing. This yeast cell divides asymmetrically, giving rise to a larger mother cell and a smaller daughter cell, leaving a circular bud scar on the mother cell's surface at the site of division. Thus, the age (counted in generations) of a mother cell can simply be determined by counting the number of bud scars on its surface. However, counting of the bud scars is labour intensive and time consuming and cannot be used as such as a screening method to isolate cells with an increased life span. Methods to isolate mutant yeasts with an increased life span have, amongst others, have been described in WO9505459 and US5874210. The latter patent describes a method to isolate a mutation which increases the number of divisions of yeast cells, comprising the labelling of the cell surface of the yeast cell with a fluorescent marker, thereby generating fluorescent yeast cells, culturing the yeast cells under conditions for

growth of yeast cells for a period of time greater than the chronological life span of the strain, selecting the fluorescent cells by fluorescence-activated cell sorting and replating the fluorescent yeast cells. However, although this method may indeed give an enrichment of strains that survive longer, there is no direct selection for strains with an increased number of divisions, and non-dividing or slower dividing cells that also survive may be selected too.

In this invention, we disclose a method for specific isolation of old yeast mother cells, with an increased number of divisions by staining the bud scar chitin with fluorescein isothiocyanate (FITC)-wheat germ agglutinin (WGA) lectin and sorting by a FACS apparatus, after initial enrichment of the mother cells through magnetic-based sorting. The process is presented in Figure 1. Said method can be used to isolate genes or mutations involved in aging.

Much attention has been focussed on the hypothesis that oxidative damage plays an important role in aging (Shan *et al.*, 2001; Hamilton *et al.*, 2001) and there is a generally accepted relation between oxidative stress and aging (Tanaka *et al.*, 2001). Moreover, mutations in genes related to protection against oxidative stress have a clear influence on life span, both in *S. cerevisiae* and *Caenorhabditis elegans* (Laun *et al.*, 2001; Ishii, 2001). This makes that the method, proposed here, is also suitable as an indirect selection for genes involved in oxidative stress. This is especially useful in cases where screening of libraries in an endogenous system is difficult or impossible, such as the screening of mammalian or plant libraries. Screening of such libraries may lead to new genes involved in protection against oxidative stress in general, but also, in case of mammalian cells, to genes involved in AAD's and/or diseases caused by oxidative stress, especially neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease and Huntington's disease (Calabrese *et al.*, 2001)

A frequently practiced strategy in searching genes responsible for aging is by selecting survivals after exposure cells to stresses. Then a question constantly remaining is whether the genes picked up are in response to the stress treatment rather than involved in aging, because of the complexity of the process. The invention described here, however, provides an alternative that allows direct hunting of genes with potential anti-aging functions from various libraries or library combinations of eukaryotic organisms. Yeast lines are selected in a more natural condition, and also with advantages of high throughput, high efficiency, and short

time investment. Obviously, this invention has a great potential for rational drug design and development of therapies and prevention in the field of age-related diseases.

- 5 It is a first aspect of the invention to provide a method to screen genes involved in aging and/or AAD's and/or oxidative stress, comprising a) mutation or transformation of a yeast cell b) cultivation of said cell c) enrichment of the population for mother cells d) labelling said mother cells with a WGA- based label and e) isolation of the highly labelled cells.
- 10 To obtain a sufficient distinction between old cells and young cells, it is essential to use a marking of the bud scars that is sufficiently linear with the number of scars, and is not or only weakly interacting with other cell wand compounds. Surprisingly we found that WGA can bind with the chitin in the bud scar, without major interference with other cell compounds, so that the amount of WGA bound is a
- 15 reliable measurement of the number of bud scars. The WGA bound is then measured using a WGA-based label. A WGA-based label, as used here, may be any kind of label that allows quantifying the amount of WGA bound to the cell and may be, as a non-limiting example, WGA coupled to a stain, or a detectable antibody that binds to WGA. Detectable antibodies are known to the person skilled
- 20 in the art and may be, as a non-limiting example, rabbit antibodies that can be detected by a labelled anti-rabbit antibody. The labelling of mother cells with a WGA based label may be a one step process, whereby labelled WGA is bound to the cell, or a two step process, whereby in a first step, WGA is bound to the bud scars, and in a second step, the bound WGA is labelled. A preferred embodiment is a method
- 25 according to the invention, whereby said WGA based label is FITC labelled WGA. Preferably, said isolation of highly stained cells is based on FACS sorting. Methods for the enrichment of the population of mother cells are known to the person skilled in the art and may be based on, as a non-limiting example, staining of the cell wall of the cells at a certain point in the growth phase, followed by continuation of the
- 30 culturing and sorting of the stained cells. Alternatively, the cells may be antibody labelled. Preferably, said enrichment of the population of mother cells is a magnetic-based sorting. Instead of being based on a global cell wall labelling as described above, the enrichment of the population of mother cells may be based on the labelling of a fraction of the mother cells, such as a bud scar based labelling. In

fact, the enrichment of the mother cells may be carried out by a first WGA based labelling and sorting, whereby the enriched mother cells are subjected to a second WGA based labelling and sorting. The labelling method in the first and second round may be different.

5 Methods to mutate yeasts are known to the person skilled in the art and include, but are not limited to chemical and physical mutagenesis, such as ethyl methane sulphonate (EMS) treatment, or UV treatment. Methods to transform yeast are also known to the person skilled in the art and include, but are not limited to protoplast transformation, lithium acetate based transformation and electroporation. The yeast
10 transformation may be carried with one or more nucleic acids, up to a complete library. The nucleic acid used is not necessarily yeast nucleic acid, but may be from any origin, as long as it is functionally expressed in yeast. Preferred examples of nucleic acids are mammalian nucleic acids, such as human nucleic acid, and plant nucleic acid, whereby said nucleic acids are cloned in a yeast expression vector.
15 Preferably, the yeast is transformed with an expression library. The nucleic acid that is transcribed into mRNA does not necessarily be translated into protein, but may exert its effect as antisense RNA. Indeed, it is an additional advantage of the method that it can detect in one screening experiment both the effect of overexpression of a protein, as well as the effect of downregulation of a protein by
20 blocking the translation of an endogenous messenger by a homologous antisense RNA, resulting from the expression library.

Another aspect of the invention is a gene or functional gene fragment isolated with the method, according to the invention. Said functional fragment may encode for a polypeptide, that directly affects aging and/or an AAD and/or oxidative stress, or it
25 may be transcribed into antisense RNA, which affect aging and/or an AAD and/or oxidative stress by silencing an endogenous gene. Preferably, said gene or functional gene fragment is selected from the nucleic acid listed in table 2. More preferably, said gene or functional gene fragment comprises a sequence as represented in SEQ ID N° 1, 3, 5, 7, 8, 9, 11, 13, 15, 16, 17, 19, 21, 22, 23, 24, 25,
30 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52 or 53. Even more preferably, said gene or gene fragment is essentially consisting of a sequence as represented in SEQ ID N° 1, 3, 5, 7, 8, 9, 11, 13, 15, 16, 17, 19, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52 or 53. Even more preferably,

said gene or functional gene fragment is consisting of a sequence as represented in SEQ ID N° 1, 3, 5, 7, 8, 9, 11, 13, 15, 16, 17, 19, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52 or 53. A preferred embodiment is a gene fragment, isolated with the method, essentially consisting of SEQ ID N° 11, preferably consisting of SEQ ID N° 11.

Another preferred embodiment is a gene fragment, isolated with the method, essentially consisting of SEQ ID N° 16, preferably consisting of SEQ ID N° 16.

Still another aspect of the invention of the use of a gene or functional gene fragment isolated with the method according to the invention to modulate aging and/or to modulate the development of AAD's and/or to protect against oxidative stress. Preferably, said modulation is an inhibition of aging. Preferably, said gene or gene fragment is selected from the nucleic acids listed in table 2. More preferably, said gene or gene fragment comprises a sequence as represented in SEQ ID N° 1, 3, 5, 7, 8, 9, 11, 13, 15, 16, 17, 19, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52 or 53. Even more preferably, said gene or gene fragment is essentially consisting of a sequence as represented in SEQ ID N° 1, 3, 5, 7, 8, 9, 11, 13, 15, 16, 17, 19, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52 or 53. Even more preferably, said gene or gene fragment is consisting of a sequence as represented in SEQ ID N° 1, 3, 5, 7, 8, 9, 11, 13, 15, 16, 17, 19, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52 or 53. A preferred embodiment is the use of a functional gene fragment, essentially consisting of SEQ ID N° 11, preferably consisting of SEQ ID N° 11. Another preferred embodiment is the use of a gene fragment, isolated with the method, essentially consisting of SEQ ID N° 16, preferably consisting of SEQ ID N° 16.

Another aspect of the invention is a polypeptide, encoded by a gene or functional gene fragment isolated with a method according to the invention. Preferably, said modulation is an inhibition of aging and/or inhibition of the development of an AAD. Preferably, said polypeptide is encoded by a nucleic acid listed in table 2. More preferably, said polypeptide is encoded by a nucleic acid comprising SEQ ID N° 1, 3, 5, 7, 8, 9, 11, 13, 15, 16, 17, 19, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52 or 53. Even more preferably, said polypeptide is encoded by a nucleic acid essentially consisting of

SEQ ID N° 1, 3, 5, 7, 8, 9, 11, 13, 15, 16, 17, 19, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52 or 53. Even more preferably, said polypeptide is encoded by a nucleic acid consisting of SEQ ID N° 1, 3, 5, 7, 8, 9, 11, 13, 15, 16, 17, 19, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52 or 53. Even more preferably, said polypeptide comprises SEQ ID N° 2, 4, 6, 10, 12, 14, 18, or 20. Even more preferably, said polypeptide is essentially consisting of SEQ ID N° 2, 4, 6, 10, 12, 14, 18 or 20. Even more preferably, said polypeptide is consisting of SEQ ID N° 2, 4, 6, 10, 12, 14, 18 or 20. A preferred embodiment is a polypeptide, essentially consisting of SEQ ID N° 12, preferably consisting of SEQ ID N° 12. Still another preferred embodiment is a polypeptide encoded by a nucleic acid essentially consisting of SEQ ID N° 16, preferably consisting of SEQ ID N° 16

Still another aspect of the invention is the use of a polypeptide, encoded by a gene or functional gene fragment, isolated with a method according to the invention, to modulate aging and/or to modulate the development of an AAD and/or to protect against oxidative stress. Preferably said modulation is an inhibition of aging and/or inhibitor of the development of an AAD. Preferably, said polypeptide is encoded by a nucleic acid selected from the nucleic acids listed in table 2. More preferably, said polypeptide is encoded by a nucleic acid comprising SEQ ID N° 1, 3, 5, 7, 8, 9, 11, 13, 15, 16, 17, 19, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52 or 53. More preferably, said polypeptide comprises SEQ ID N° 2, 4, 6, 10, 12, 14, 18 or 20. Even more preferably, said polypeptide is essentially consisting of SEQ ID N° 2, 4, 6, 10, 12, 14, 18 or 20.

Most preferably, said polypeptide is consisting of SEQ ID N° 2, 4, 6, 10, 12, 14, 18 or 20. A preferred embodiment is the use of a polypeptide, essentially consisting of SEQ ID N° 12, preferably consisting of SEQ ID N° 12, to modulate aging and/or to modulate the development the development of an AAD. Preferably, said modulation is an inhibition of aging and/or an inhibition of the development the development of an AAD. Still another preferred embodiment is the use of a polypeptide, encoded by a nucleic acid comprising SEQ ID N° 16, preferably essentially consisting of SEQ ID N° 16, more preferably consisting of SEQ ID N° 16, to modulate aging and/or to modulate the development the development of an AAD.

Still another aspect of the invention is the use of an antisense RNA encoded by a gene or a functional gene fragment, isolated with a method according to the invention, to modulate aging and/or to modulate the development the development of an AAD. In such an application, the gene or functional gene fragment is operationally linked to a promoter, in such a way that an antisense RNA, complementary to the mRNA encoding the polypeptide normally encoded by said gene or gene fragment, is transcribed. Preferably, said gene or functional gene fragment encoding the antisense RNA comprises SEQ ID N° 7, 8 or 15. Even more preferably, said modulation of aging is an inhibition of aging and/or an inhibition of the development the development of an AAD.

Definitions

Gene as used here refers to a region of DNA that is transcribed into RNA, and subsequently preferentially, but not necessarily, translated into a polypeptide. The term is not limited to the coding sequence. The term refers to any nucleic acid comprising said region, with or without the exon sequences, and includes, but is not limited to genomic DNA, cDNA and messenger RNA. As, on the base of these sequences, it is evident for the person skilled in the art to isolate the promoter region, the term gene may include the promoter region when it refers to genomic DNA.

Nucleic acid as used herein refers to a polymeric form of nucleotides of any length, either ribonucleotides or deoxyribonucleotides. This term refers only to the primary structure of the molecule. Thus, this term includes double- and single-stranded DNA, and RNA. It also includes known types of modifications, for example, methylation, "caps" substitution of one or more of the naturally occurring nucleotides with an analog.

Functional fragment of a gene involved in aging is every fragment that, when tested with the method according to the invention, still gives a positive response. Typically, functional fragment are fragments that have deletions in the 5' and/or 3' untranslated regions. Alternatively, the functional fragment may be an antisense fragment, encoding an RNA that is silencing an endogenous gene, or functions as RNAi. As the coding sequence on its own is also considered as a functional fragment, as it is evident for the person skilled in the art that it may be functional when it is placed between suitable heterologous 5' and 3' untranslated sequences.

Polypeptide refers to a polymer of amino acids and does not refer to a specific length of the molecule. This term also includes post-translational modifications of the polypeptide, such as glycosylation, phosphorylation and acetylation.

5 *Aging* as used here includes all forms of aging, particularly also aging-associated diseases (AAD's). AAD's are known to the person skilled in the art and include, but are not limited to arteriosclerosis, Parkinson's disease and Alzheimer's disease.

Brief description of the figures

10 Figure 1. Scheme of the bud scar sorting (BSS) system for yeast M-cells. The BSS system contains two major steps. The first step at the left side of the figure, magnetic sorting of biotinylated M-cells and re-growth of sorted M-cells to desired generations when needed. The second step at the right side of the figure, WGA staining of bud scars and sorting of longer life M-cells according to bud scar staining.

15 Figure 2. Flow cytometric assay of yeast cells labelled with WGA-FITC and streptavidin-PE.

Yeast cells (M-cell) are grown for 5 to 6 generations (G5-6) after biotin labelling, sorted via MACS, and then simultaneously labelled with WGA-FITC and streptavidin-PE. A: shows a clear separation of the PE red-fluorescent mother cells (gated M-cell) from the non-PE fluorescent daughter cells (gated D-cell). B: hardly detects the PE fluorescent signal in the depleted daughter cells. C and D: the layout of FSC versus SSC, the gated M-cells mainly appeared at higher FSC/SSC values representing a large cell size population (C) compared to a small cell size population of D-cells at lower FSC/SSC values (D). E and F: the M-cell population gives strong WGA-FITC staining (E) than the D-cell population (F).

20

25

Figure 3. Bud scar staining of yeast cells. INVSc-1 cells (M-cells) were biotinylated and cultured in SD medium. M-cells at G5-6 were magnetically sorted. Staining of bud scars with WGA-FITC was revealed with a Zeiss LSM410 confocal microscope.

30

Figure 4. Screen of a human cDNA library via FACS.

A cDNA library from HepG2 hepatoma cells was transformed into the yeast strain INVSc-1 (pEX2) (See Materials and Methods). The transformed yeast population was first labeled with biotin and then cultured in S-glycerol medium. The initial biotinylated M-cells of approximately G14 (14 generations) were obtained by running

35

two magnetic sorting and regrowth cycles, and were then double labelled with WGA-FITC and streptavidin-PE. The older mother cells were gated according to PE staining and big cell size which represented as high FSC (A). Flow sorted older mother cells (gate Old-M) show a strong WGA-FITC signal (B).

5

Figure 5. Flow cytometric dead cell assay using PI staining.

Flow cytometric analysis of cell death using PI staining was performed in a ferritin L chain clone (pEX2-FL) and its parent line of INVSc-1 (pEX2). Yeast cells were grown up to 6 generations. The gate R1 was set around PI-positive cells that cover the dead cells, the gate R2 around the PE-positive cells that represents the M-cell population and the gate R3 around the D-cell population. In the panel A, it shows 16,3% dead cells for the ferritin L chain clone. In panel B, a 33% dead cell was observed in the control line.

10

15

Figure 6. Resistance of ferritin containing yeast to H₂O₂ (1mM) stress.

Cells transformed with the plasmids as indicated were exponentially grown at 30° C to an OD₆₀₀ of approximately 0.5. Cells were treated with 1mM H₂O₂ during various times. Samples were diluted and plated on YPD solid media to monitor cell viability. Cl2-ferritin indicates the cell line containing the ferritin-fragment expression vector of pGAL10-FL. Its parent line transformed with the empty vector of pSCGAL10-SN was used as control.

20

Figure 7. Life span of *C. elegans* carrying the human Ferritin Light Chain (*FTL*) gene. Animals were injected with a L4759 plasmid containing human *FTL* gene. Controls were injected with empty plasmids. pRF4 containing the dominant phenotypic marker *rol-6(su1006)* was coinjected in both cases. Results are cumulative from four independent experiments with more than 25 animals per trial. Life-span is defined as the day when the first transformed larvae hatched until their death. Animals carrying copies of the human *FTL* gene lived significantly longer (13.54 ± 0.269 days) than controls (12.50 ± 0.266 days).

25

30

Figure 8. Study of the aging phenotype of yeast $\Delta fob1$ strain by the mixed-growth system. A mixture of $\Delta fob1$ strain and parent BY4742, were biotinylated and grown in SD medium as described in example 7. G20 (the point after 20 generations) was

obtained by running three cycles of magnetic sorting and regrowth. The results show an increased frequency of $\Delta fob1$ cycling M-cells at G20, illustrating a longer life span.

Figure 9. Comparison of the viability of *FTL* strain with its parents.

5 The initial mixture of M-cells (*FTL* and INVSc-1) was biotinylated and grown in minimal SD and S-glycerol media as described in materials and methods. The ratio of viable M-cells in the mixture at different ages was determined by plating. Data for cells grown in the *FTL* gene inducing S-glycerol medium, are presented at the right side of the figure, while data for the control are shown on the left side, indicating that
10 the difference in aging is clearly due to the ferritin expression. In a separate experiment, doubling times of both strains were carefully tested and found to be equal.

Figure 10. Ferritin L prevents fast aging in presence of iron in yeast as tested by
15 micromanipulator experiment

Life spans of human partial ferritin and full ferritin transformed in strain BY4741. S-raffinose was used as carbon source for inducing expression of ferritin. An excess of iron was added in the medium with 500 μ M FAC and 80 μ M ferrichrome. At least 60 cells were included in each of three life span assays. Both partial and full ferritin had
20 a longer average life span (17.85 G and 15.58G) than the control (12.19).

Examples

Materials and methods to the examples

25 *Strains and Media*

The following *S. cerevisiae* strains were used: INVSc-1 (Invitrogen, San Diego, CA); BY4741 and BY4742 (Euroscarf, Frankfurt, Germany) as well as the BY4742-derived $\Delta fob1$ strain (Euroscarf; accession No. Y14044). Strains were grown at 30°C in rich YPD medium (2% dextrose, 2% bactopectone and 1% yeast extract) or minimal SD
30 medium (0.67% yeast nitrogen base without amino acids, 2% dextrose and 0.077% complete supplement mixture - uracil). The INVSc-1 and BY4741 strains used for library screening were grown in S-glycerol, S-galactose or S-raffinose media, where dextrose is replaced with 3% glycerol, 2% galactose or 2% raffinose, respectively. S-glycerol was used to induce expression of genes cloned in pEX2, whereas S-

galactose was used to induce expression of genes cloned in pSCGAL10-SN. Media were solidified with 2% agar.

Cloning and overexpression of a human cDNA library

5 To recover mRNA from various responses, a pool of equal proportions of human HEPG2 cells, subjected to different treatments, was used for library construction. These treatments included heat shock for 1.5 h at 42.5°C, 1 mM dithiothreitol, 100 U/ml interleukin-6 and 10^{-7} M dexamethasone. Construction of cDNA libraries was carried out essentially as described previously (Declercq et al. 2000). cDNA was
10 cloned at the site of SfiI/NotI in the vectors pEX2 (BCCM/LMBP Plasmid Collection, Ghent University, Belgium; accession No. 2890) and pSCGAL10-SN (BCCM/LMBP Plasmid Collection, accession No. 2471). cDNA expression is driven by the cytochrome c promoter in pEX2 and by the GAL10 promoter in pSCGAL10-SN. Yeast strain INVSc-1 was used as the host for pEX2 library transformation. The pSCGAL10-SN library was transformed to the BY4741 strain. Transformations were performed as
15 described previously (Gietz and Woods, 2001). Approximately 3.5×10^5 colonies from each transformation were produced.

Magnetic sorter based preparation of yeast mother cells (M-cell)

20 Cells were cultured at 30°C in liquid medium, such as minimal SD medium or in the specific induction medium, to OD₆₀₀ of 0.7-1 and were collected by centrifugation. All cells harvested were used as M-cells. The biotin labelling of M-cells was carried out essentially as described previously (Smeal et al., 1996). Before labelling, M-cells were washed twice with cold phosphate-buffered saline (PBS; pH 8.0), resuspended in PBS
25 to a concentration of 2.5×10^7 cells/ml and then incubated with 0.1 mg/ml Sulfo-NHS-LC-Biotin (Pierce Chemical Company, Rockford, IL) for 30 min at room temperature under gentle shaking. The free biotin reagent was removed by two washings with PBS. Biotinylated M-cells were grown in liquid medium for a desired number of generations (up to G7 in our conditions; culture was not allowed to exceed OD₆₀₀ = 1).

30 The separation of mother cells from the daughter cells they produced was carried out via magnetic cell sorting. This was realized by coupling the biotinylated mother cells to magnet beads by incubating 10^7 mother cells with 80 µl of Anti-Biotin MicroBead (Miltenyi Biotec, Germany) in 1ml PBS pH 7.2 for 1hour at 4°C. Unbound

beads were removed by washing twice with PBS. M-cells were isolated with a magnetic sorter according to the supplier's protocol (Miltenyi Biotec). When needed, these sorted M-cells can be further grown in liquid medium for additional generations and isolated again by the magnetic sorting system.

- 5 The purity of sorted mother cells was determined on the basis of streptavidin binding. About 10^7 biotinylated cells were stained with 3 μ g streptavidin-conjugated R-phycoerthrin (PE) (Molecular Probes) in 1 ml of PBS pH 7.2 for 1 hour at room temperature in total darkness. Then cells were washed twice with PBS and suspended in 2 ml of PBS pH 7.2. The yeast cells with more bud scars were
10 recognised as a high intensity of FITC signals.

WGA-based bud scar staining

- The bud scars of yeast cells were stained with fluorescein isothiocyanate (FITC)-labelled WGA lectin (Sigma). The staining was carried out by adding 10^7 yeast cells
15 together with 12 μ g WGA-FITC in 1 ml of PBS pH 7.2 for 1.5 hours at room temperature, in the dark. After two washing steps with PBS to remove the free WGA-FITC reagent, yeast cells were resuspended with PBS to a concentration of 0.5×10^7 cell/ml for FACS analysis.

Propidium iodide (PI) staining

- PI (Sigma) was freshly dissolved in PBS buffer to a final concentration of 1mg/ml as stock solution. For staining, yeast cells were suspended in PBS pH 7.2 to approximately 10^7 cell/ml and then, 3 μ l of PI stock solution was added into 1 ml yeast cell suspension. The sample was run within 5-10 minutes on a flow cytometer
25 (Becton Dickinson), which is capable of measuring red fluorescence (with a band pass filter >650). No washing steps were included.

Set-up of Becton Dickinson FACScan

- Analysis of FITC, PE and PI labelling of the cell population was accomplished at an
30 excitation wavelength of 488 nm, using a 15 mWatt argon ion laser. FITC emission was measured as a green signal (530 nm peak fluorescence) by the FL1 detector, PE was measured as an orange signal (575 nm peak fluorescence) by the FL2 detector, and PI was measured as a red signal (670 nm peak fluorescence) by the FL3 detector. The FACScan flow cytometer (Becton Dickinson) was operated

according to the standard protocol of the supplier. For multi-colour staining, electronic compensation was used among the fluorescence channels to remove residual spectral overlap. A minimum of 10,000 events was collected on each sample. Analysis of the multivariate data was performed with CELLQuest software
5 (Becton Dickinson Immunocytometry System).

Transformation and aging assay in nematode

The expression vector of human ferritin fragment (FTL) for *C. elegans* was derived from L4759 by replacing the GFP with FTL fragment.

10 Wild-type *C. elegans* strain (N2) was used as host for FTL expression. The animals were cultured and handled as described (Brenner, 1974). The transient overexpression of human FTL was carried out according to Jin (1999) using an Eppendorf FemtoJet-TransferMan NK injection system (Eppendorf, Leuven, Belgium). 25-30 worms were injected with plasmid carrying the human FTL gene or
15 control plasmid. Plasmid pRF4, which carries the dominant rol-6(su1006) allele was coinjected to mark transformed progeny. After a one-hour recovery period in M9 buffer, injected animals were allowed to lay eggs for approx. 40 hours on plates containing nematode growth medium (NGM) and a lawn of *E. coli* bacteria (OP50) as food. Transformed eggs were predominantly laid during the last 20 hours
20 resulting in a fairly synchronous experimental cohort. Subsequently, the injected animals were removed and progeny (F1) was allowed to grow at 24°C. Fourth stage larvae or young adults showing the Roller phenotype were transferred onto separate plates (NGM + OP50) containing 300 µM 5-fluoro-2'-deoxyuridine (FUDR, Sigma) to prevent progeny (F2) production. Live/dead scoring was carried out daily. Lifespan
25 is defined as the day when the first transformed larvae hatched until their death.

Construction of a full ferritin clone

A ferritin PCR fragment (end to stop codon) was generated from the hepatoma cDNA library by using specific primers (5'-ctacgagcgtctcctgaagatgc3' and
30 5'-cgcggatccaagtcgctgggctcagaaggctc-3'). This fragment was cloned directly into the TOPO vector (Invitrogen, The Netherlands) and then digested with NotI, generating a NotI fragment. Subsequently, the NotI fragment was inserted in the NotI site of ferritin light fragment clone (pGAL10-FL), resulting a 750 bp full ferritin clone in pSCGal-SN-10.

35

Example 1: Magnetic based sorting of yeast M-cells

To use yeast as an aging model, the first step needed is the development of a system, which allows the isolation of a relatively pure population of old yeast cells. The method for distinguishing and separation of *S. cerevisiae* cells between
5 generations is based on the fact that daughter cells have a wall that is newly formed and do not have any detectable wall remnants of the mother cells. Cells from an overnight culture of *S. cerevisiae* strain INVSc1 in minimal SD medium were covalently coated with biotin and designated as mother cells (M-cell). The M-cells were inoculated into fresh medium, and allowed to grow for 5-6 generations as
10 determined by the cell density that is measured by a UV-visible spectrophotometer (Shimadzu). After loading with anti-biotin beads, M-cells were sorted out using a magnetic sorter or MACS (Materials and Methods).

The purity of the collected M-cells was determined by staining with streptavidin-PE, which specifically binds to biotin coated on the cell wall of M-cells, followed by flow
15 cytometric analysis. Due to the reaction of biotin with streptavidin-PE, high density staining of biotinylated M-cells was shown. As show in Figure 2A, there was clear separation between stained M-cells and unstained daughter cells (D-cell) populations. Gate and marker were positioned to exclude D-cells from the M-cell population. In the layout of FSC versus SSC, as the matter of fact, the gated M-cells
20 mainly appeared at high FSC/SSC values representing a large cell size population (Fig. 2C) compared to a small cell population of D-cells which mainly located at lower FSC/SSC values (Fig. 2D). Statistic analysis showed that the purity of the isolated M-cells reached more than 85%. Figure 2B shows a PE staining performed on a depleted D-cell population, which hardly shows any positive signal.

25

Example 2: WGA based staining for analysis of yeast life span

Wheat germ agglutinin (WGA, *Triticum vulgare*) is the first lectin of which the amino acid sequence was completely determined (Wright, 1984). WGA is a mixture of several isolectins (Rice and Etzler, 1975). Sharing similar carbohydrate binding
30 properties with other lectins, WGA reacts strongly with the chitobiose core of asparagines linked oligosaccharides, especially with the Man β (1,4)GlcNAc β (1,4)GlcNAc trisaccharide (Yamamoto et al., 1981).

One of the most striking features of the cell surface during aging *S. cerevisiae* is the accumulation of chitin-containing bud scars. To verify whether WGA can be used for

specific labelling of chitin in yeast bud scars, the yeast strain INVSc-1(pEX2) was incubated with the FITC-conjugated WGA. The enriched, magnetically sorted M-cells were subjected to WGA reaction.

Under a fluorescence microscope we found that the major part of the fluorescent signal for WGA-FITC staining was co-localizing with the bud scar rings (Fig. 3). Moreover, the number of stained bud scars (6 bud scars) was consistent with the expected age of the M-cells as estimated by cell density measurement of the culture (5-6 generation). This observation demonstrated that, under the conditions used, WGA is specifically binding to the chitin of bud scars and hardly gives any fluorescence, caused by binding to compounds in the normal cell wall. Therefore, the possibility was examined to use WGA as a tool to stain bud scar for analysis of yeast life span. The isolated M-cells and depleted D-cells (as seen in Fig. 2C and 2D) were simultaneously stained with streptavidin-PE and WGA-FITC. As shown in Figure 2E-2F, D-cells that were negative for streptavidin-PE staining showed low FITC signal (Fig. 2F), whereas M-cells, which were positive in streptavidin-PE staining, showed a much stronger FITC staining (Fig. 2E). Under the fluorescent microscope, we observed that most M-cells contained 5-6 bud scar rings, which were strongly labelled by WGA-FITC, while most D-cells had only 1-2 bud scar rings. This observation indicated that there was a good linear correlation between the number of bud scars and the intensity of fluorescence. Therefore, it was assumed that WGA could be used as a tool for bud scar-specific staining in budding yeast cells.

Example 3: Application of using WGA to screen a human cDNA library

It has been reported that overexpression of certain human genes in yeast might have an influence in the frequency distribution of the yeast population (Gershon and Gershon, 2000). This overexpression of a single gene, which modulates the longevity in a single-cell system, has opened up the field of aging study to the power of yeast genetics. To screen human genes that might be involved in aging processes, a cDNA library from hepatoma cells was constructed and transferred into the yeast strain INVSc-1(pEX2) (See Materials and Methods). The transformed yeast population was first labelled with biotin and then cultured in a Bioreactor (AppliTek), for about 14 generations, as deduced from the cell density. According to the method described above, the initial biotinylated M-cells were isolated by

magnetic beads described herein and then labelled with WGA-FITC. By flow cytometric analysis (Fig. 4A), the M-cell population had a high density of WGA-FITC staining (gate M-cell), whereas D-cells showed a lower fluorescent staining (gate D-cell). As shown in Figure 4, older M-cells, gated as Old-M population, which were supposed to have a longer life span, were marked on high FITC intensity combined with high FSC, and then were flow sorted by FACS. From 9 colonies, the gene, overexpressed in the yeast cell was sequenced, and the results are summarized in Table 1. The growth rate was tested by measuring the doubling time of each strain in the liquid medium. The result showed that the growth rate of all 9 clones as well as the parent line were similar.

One of the colonies contained a gene fragment encoding ferritin light (FL) chain (M1147.1; Af119897.1). To verify whether the overexpression of this gene could influence the life-span of the yeast cell or not, an analysis of cell death using PI staining was performed in this ferritin L chain clone (CI2-FL) using its parent line of INVSc-1(pEX2) as a control. Ten million M-cells for each cell line were isolated. As shown in Figure 5, on the FSC versus PE (FL2) dot plot, a gate R2 was set around the PE-positive cells that represents the M-cell population while a gate R3 was set around the D-cell population. At the same time, on the FSC versus PI (FL3) dot plot, a gate R1 was set around PI-positive cells that cover the dead cells. As seen in Figure 5, cell death in culture occurred mainly in the M-cell population, but was barely detected in the D-cell population. Statistical analysis for dead cells (PI-positive) showed a higher frequency in control cells (33% death) compared to that in CI2-FL cells (16.3% death). This result indicates that over expression of human ferritin L chain in yeast cells prevents early cell death.

Example 4: additional screening experiments

To confirm the usefulness of the method, additional screening experiments were set up, using the same outline as described above both using the pEX2 library and the pSCGAL10-SN library. The results of the additional screening experiments are listed in Table 2, and identified by their genbank accession number. Several results of the first screening have been confirmed, illustrating the usefulness and the reliability of the method.

Example 5: protective effect of the ferritin fragment on hydrogen peroxide treatment

One of the colonies contained a gene fragment encoding ferritin light (FTL) chain (M1147.1; Af119897.1) cloned in pSCGAL10-SN. The plasmid was indicated as pGAL10-FL. Ferritin is ubiquitously distributed in the animal kingdom. It is composed of two subunits, the heavy chain (H) and the light chain (L). Ferritin plays a major role in the regulation of intracellular iron storage and homeostasis. One of the functions is to limit iron availability for participation in reactions that produce free oxygen radicals, which have the potential to damage lipids, proteins and DNA. Indeed, several reports have implicated that ferritin is involved in the protection against oxidative stress, such as stress induced by hydrogen peroxide. However, there is not such ferritin-like protein present in yeast, and anti-oxidative activity of ferritin fragments was never demonstrated. To test whether the human ferritin fragment plays a role as an antioxidant in yeast, we examined the partial-ferritin L clone (Cl2-ferritin), which was isolated by the method according to the invention, against H₂O₂ stress.

The condition for treatment of the cells was essentially the same as described by Jamieson et al. (1994). Exponential phase cultures of strain BY4741 that contained the empty vector pSCGAL10-SN (Control) and the ferritin expression vector (FTL – indicated as Cl2-ferritin) respectively, were grown aerobically in S-galactose medium at 30° C. The cell cultures were then challenged to a lethal concentration of H₂O₂ (1mM). Cell survival was monitored by taking samples at 0, 30 and 60 min, diluting the samples in the same medium and plating aliquots on YPD plates.

The experiment showed that, compared with control line, ferritin cells are significantly more resistant to treatment with 1mM H₂O₂ (Fig. 6).

Example 6: Transgenic nematode overexpressing the Ferritin Light chain

Although on the cellular level, there might be some conserved mechanism of aging processes throughout evolution (Martin et al., 1996), it is easy to imagine that in different species some underlying distinctive ways of intercellular regulation also contribute to reach their fate (Guarente 2001). In this sense, results from other organisms may provide a closer vision on the postulated function of human *FTL* gene involved in aging. Therefore, we tested whether FTL might affect lifespan in *C. elegans*, a multicellular organism, too. Indeed, as shown in Figure 5, animals

carrying human *FTL* genes appeared an average life of 13.5 days, which is 8% longer than the control line and statistically significant ($p=0.006$, two-way ANOVA). Many reports in *C.elegans*, *Drosophila* and mice are consistent with the hypothesis that oxidative damage accelerates aging, and that increased resistance to oxidative damage can extend lifespan (Finkel and Holbrook, 2000). The consistency that the expression/overexpression of human *FTL* gene was in favour in extending the lifespan in mono-cellular yeast and multi-cellular nematode supports the postulation that ferritin extends lifespan in cells, probably by protecting cells from oxidative stress, in a wide range of species.

A frequently practiced strategy in searching gene responsible for aging is by selecting survivals after exposure cells to stresses. Then a question constantly existing is that the genes picked up might be in response to the stress treatment rather than involved in aging, because of the complicity of the process. The screening method described here, however, provides an alternative that allows direct hunting of genes with potential anti-aging functions from various libraries or library combinations of eukaryotics. Yeast lines are selected in a more native condition, and also with advantages of high throughput, high efficiency, and short time consuming. Obviously, it has a great potential in application in rational drug design and therapies development in the field of age-related diseases preventing / treatments.

Example 7: elaboration of the mixed culture experiments

Based on the fact that a parental yeast strain and its direct derivative have a similar cell cycle rate, a mixed culture method has been developed to verify the long-living character of a transformed yeast strain when these strains are grown together in the same culture.

Two (or possibly more than two) yeast strains with a similar growth rate are initially mixed in the same culture in an equal ration (50% each in the case of two strains).

The strains can be distinguished from each other by the use of a selective marker.

The initial inoculated cells, called mother cells (M-cell), are labelled with biotin, and are grown together in the same culture during their entire life span. Mother cells at different generation points are sampled and collected by a magnetic system (MACS), similar to the method described in example 1. The ratio of living M-cells from the two strains is determined by the use of the selective marker. If the two strains have the

similar lifespan, the ratio of two viable strains will stay the same at different generation time points; otherwise, the ratio will change. This method is essentially based on the screening method, whereby the identification of the long living cells is not carried out by WGA staining, but by direct count of the number of living mother cells of the transformed strain(s), compared to the number of living mother cells of the parental strain.

FOB1 is required for the replication fork block. A *FOB1* mutation results in a decreased rDNA recombination rate and an increase in yeast life-span of 70%. The growth rate of the $\Delta fob1$ mutant strain, as measured, is similar to its parental strain.

Therefore, the long-living $\Delta fob1$ strain with its parental strain BY4742 was used to develop the mixed-growth system.

The initial mother cells were prepared as follows: a first pre-culture was made by inoculating BY4742 and $\Delta fob1$ cells (from freshly grown on a SD plate) in 5 ml of SD medium, respectively. The culture was incubated at 30 °C on a shaker at 250-300 rpm overnight. A second pre-culture was made by inoculating the first pre-culture into 5 ml of SD medium at a cell density of $OD_{600} = 0.001 \sim 0.005$. These cells were incubated until the culture reached a cell density of $OD_{600} = 0.5 \sim 0.7$. Cells were collected by centrifugation of the culture at 4 °C for 5 min at 3000 rpm. The cell pellet was washed twice with pre-cooled PBS (pH 8) and resuspended in PBS at a cell density of $OD_{600} = 5$ (approximately 5×10^7 cells/ml). The biotinylation of cells was performed in an eppendorf tube, in 1 ml reaction volume consisting of 0.5 ml of above-mentioned cells (2.5×10^7 cells) and 0.5 ml of 1 mg/ml biotin (Sulfo-NHS-LC-Biotin). The mixture was incubated for 30 min at room temperature with a gentle shaking. The biotinylated cells were centrifuged for 5 min at 13000 rpm and washed twice with 1 ml of cold PBS to get rid of free biotin. These cells were used as initial mother cells (M-cell).

A 100ml mixed-growth culture of BY4742 and $\Delta fob1$ was set up by inoculating 1×10^7 biotinylated M-cells from each strain (mother cells) at ratio of 1:1 in a SD medium. The mixed-growth culture was incubated at 30 °C on a shaker at 250-300 rpm. The culture density was not allowed to exceed $OD_{600} > 1$.

After growing several generations (up to 7-generation in our condition), the M-cells were labelled with anti-biotin microbeads and isolated using the magnetic system (MACS). The purity of M-cells was determined by FACS (fluorescence-activated cell sorter) after staining M-cells with streptavidin-conjugated with PE. Using these

conditions, more than 90% M-cells could be obtained. After the final magnetic sorting, the ratio of viable M-cells was measured.

Mixed M-cells samples were plated at about 500 cells per plate on YPD and YPD/geneticin plates to determine the ratio of mother cells of the two strains at different generation points. Plates were incubated for three days at 30 °C. The ratio of BY4742 and $\Delta fob1$ mother cells was monitored by counting the colonies on the two kind of plates. The total viable number of M-cells could be determined on the YPD plate, while the number of viable $\Delta fob1$ M-cells could be derived from YPD/geneticin plate.

As shown in Figure 8, the mixed M-cell group had similar amounts of the two strains at G0, while at G20 M-cells from $\Delta fob1$ were dominant (96%) among the cells sorted and collected with the magnetic sorting system. This result confirms that the mixed-growth method could indeed be used to distinguish the longer living yeast strain from its control.

Example 8: Confirmation of aging phenotype of ferritin strain by mixed-growth system

A kinetic analysis for growth rate of the ferritin yeast (FTL) and its parental strain INVSc-1 (with a geneticin-selectable marker) revealed a similar rate. About an equal amount of two strains was mixed, as described above, but using S-glycerol medium to obtain induction of the ferritin expression. This mixed culture was subjected to a mixed-growth experiment for determining their life span differences. After examination of the longevity of a mixed-growth of these two cell types by mixed-growth system and subsequent plating, we found that the ferritin line was predominant in the viable M-cell group after a growth of 10 generations (Figure 9). Growth of a mixture of these two lines in SD medium, in which the expression of ferritin not induced, revealed a constant viable FTL/INVSc-1 ratio. This indicates that the extended longevity of the FTL strain, compared to the age-matched INVSc-1 strain, is caused by the expression of human FTL.

Example 9: independent confirmation of effect on life span by ferritin

Iron is an essential nutrient for virtually every organism because is required as an essential cofactor for many proteins. However, excess iron can generate via the Fenton reaction highly toxic-free radicals generating oxidative damage to the cell.

Thus, cellular iron concentration must be tightly controlled. To exam whether expression of human ferritin in yeast could protect cell death upon excess iron, the lifespan analysis of ferritin strains was carried out by micromanipulaor as described previously (Kennedy et al., 1994) with the following slight modifications. Cells were
 5 pregrown on non-inducing SD medium (2% glucose), shifted to inducing S-raffinose (2% raffinose) medium with 500 μ M ferric ammonium citrate (FAC) and 80 μ M ferrichrome (Sigma), and grown for at least two generations. Cells were taken from this logarithmically growing liquid culture and transferred at low density on S-
 10 raffinose with 500 μ M FAC and 80 μ M ferrichrome plate (2% agar). The cells were then incubated at 30 °C overnight. Virgin daughters cells were isolated as buds from populations by micromanipulator and used as the starting mother cells for life span analysis. For each successive bud removed from these mother cells, they were counted one generation older. Cells were grown at 30°C during the day and at cold room overnight. Each experiment consists of at least 60 cells. The statistical
 15 analysis of life span was carried by a Wilcoxin 's test. The life span of full ferritin and partial ferritin yeast strains were significantly extended by 10 to 15% compared to their parent strain BY4741 (Figure 10). This result confirms that human ferritin light chain prevents fast aging in presence of iron in yeast.

20 **Table 1:** results of the screening of 9 positive clones

Clone number	Insert length (approx.)	Identification (Based on homology)	SEQ ID N°
1	1.6 kb	Humanin	1
2	883 bp	APOA1	3
3	1 kb	Ribosomal protein P0	5
4	2.8 kb	glutamyl tRNA synthetase	7 ⁽¹⁾
5	2.4 kb	GRSF-1	8 ⁽¹⁾
7	700 bp	ALDH1	9
8	416 bp	ferritin light chain	11
9	1 kb	Ribosomal protein S2	13
12	500 bp	Histone H2A	15 ⁽¹⁾

(1) antisense

Table 2: Results of further screening experiments. The results are grouped in mitochondrial functions, ribosomal proteins, other genes with known function, unknown functions and chromosomal fragments. The results of the first screening are not repeated in this table; however, several genes, like the ferritin fragment, have been identified in more than one screening experiment. The sequences are identified by their genbank accession number. The length of the isolated fragment may differ from the genbank sequence, and is normally shorter. Where relevant, the fragment is indicated, using the nucleotides numbers of the genbank sequence.

Mitochondrion

clone name	Function	accession number	orientation
1E3/6D8	ATP synthase 6 mRNA,	AF368271	sense
2C10	mitochondrial ATP synthase subunit 9, P3 gene copy, mRNA, nuclear gene encoding mitochondrial protein	U09813	sense
5D9	ATP synthase, H ⁺ transporting, mitochondrial F1, complex, O subunit (oligomycin sensitivity conferring protein), (ATP5O)	NM_001697 89-745	sense
9B11	ADP/ATP translocase mRNA, 3' end	J03591	sense
4D7/7H1/12D3 /13E9	NADH dehydrogenase 1	BC009316 380-685; 10-684; 138-645; 10-490	sense
6C11	NADH dehydrogenase 1	BC009316	sense
7E11	NADH dehydrogenase subunit 5 (MTND5) mRNA, RNA 4, complete cds; mitochondrial gene for mitochondrial product	AF339086	sense
10G3	mitochondrion cytochrome b gene, partial cds	U09500	sense
12F1	cytochrome c oxidase subunit III gene, mitochondrial gene encoding mitochondrial protein, partial cds	AF004341	sense
2A7	ubiquinol-cytochrome c reductase core protein II	BC003136; 763-1331	sense
1B12	monocyte chemotactic protein-3 (MCP-3)	X72308	
1F9/12H12	Wnt-13 ; metochrondrial DNA	Z71621; 1-348; 12-372	sense
7C1	12S ribosomal RNA gene, partial sequence; and tRNA-Val gene, complete sequence; mitochondrial genes for mitochondrial products	AY012136	sense

11B7	MRPS16 mRNA for mitochondrial ribosomal protein S16	AB049948.	sense
10F3/ 14H4/14H5 /7B10	clone IMAGE:5581122, mRNA; haplotype N1b mitochondrion	<u>BC035832.1</u> <u>AF381999</u> ; <u>228-726; 330-953 ; 134-</u> <u>1028 2059-2658</u>	sense

Ribosome

1S_3	ribosomal protein P0	BC005863	sense
3A5	ribosomal protein, large, P1	NM_001003.2	sense
12E12	ribosomal protein L12 (RPL12)	NM_000976	sense
6F8	ribosomal protein L14	BC029036	sense
1D12	ribosomal protein L31 (RPL31)	NM_000993	sense
1S_9	ribosomal protein S2	NM_002952	sense
6D6	ribosomal S3 (RPS3);	NM_001005.2	sense
3D1/4G10	ribosomal protein S3A	<u>BC030161</u>	sense
	v-fos transformation effector protein (Fte-1)	M84711	sense
3B9	ribosomal protein S4, X-linked (RPS4X)	<u>NM_001007</u>	sense
	scar protein	M22146	sense
4H2	ribosomal protein S4, Y-linked (RPS4Y)	<u>NM_001008</u>	sense
10E8	ribosome protein S5	BC018151	sense
14G6	ribosomal protein S6 (RPS6)	NM_001010.2	sense
4C5	ribosomal protein S10,	<u>BC005012</u>	sense
4B5/2A3	ribosomal protein S11	BC016378	sense
	Mus musculus RAD21 homolog (S. pombe) (Rad21)	NM_009009	sense
11E4	ribosome protein S16	nm_001020	sense
2E6	ribosomal protein S17 mRNA	<u>M13932</u>	sense
1D1	ribosomal protein S25	<u>BC004986</u>	sense
1C11	Wilm's tumor-related protein (QM) mRNA; RPL10	M64241	sense

Other genes from the 4th screen (pEX2 library)**Unknown functions**

2H4	likely ortholog of mouse gene rich cluster, C8	<u>NM_031299.2</u> 346- <u>end</u>	sense
3C2	clone FLC0593	<u>AF113701</u>	sense
4C11	similar to putative, clone MGC:33177 IMAGE:4823662	BC028387; 1905-end	sense
4D10	full length insert cDNA clone ZE03C06	<u>AF086514</u>	sense
4E9	hypothetical protein dJ465N24.2.1 (DJ465N24.2.1)	NM_020317; 874-1431	sense
6F6	Similar to RIKEN cDNA 1110012M11 gene	BC007883	sense

6H8	cDNA FLJ31039 fis, clone HSYRA2000221	AK055601; 1869-end	sense
7F6	cDNA FLJ13305 fis	AK023367	sense
8C10	hypothetical protein FLJ23018 (FLJ23018)	NM_024810	sense
9F4	Similar to hypothetical protein FLJ10751	<u>BC024001; 3-end</u>	sense
9G10	hypothetical protein BC013073 (LOC92703)	NM_138391	sense
10G6	similar to C50F4.16.p (LOC256281)	XM_170755	antisense
12E6	hypothetical protein MGC955	NM_024097.1	sense
14D4	clone IMAGE; 4778940 mRNA	<u>BC031919.1; 3-end</u>	sense
5S-15/114	cDNA DKFZp434O159	AL133593	sense
5S-21/57	hypothetical protein FLJ10081	NM_017991	sense
7F11	cDNA FLJ38528 fis	AK095847	sense
11H3	cDNA FLJ14279 fis	AK024341; 362-end	sense
5C4	Similar to KIAA0674 protein	BC026048	sense
2E2	cDNA FLJ14385 fis, clone HEMBA1002212, weakly similar to TYROSINE-PROTEIN KINASE 2	AK027291; 3-end	sense
7G6	KIAA0776 protein (KIAA0776)	NM_015323; 3-end	sense

Chromosome DNA seq.

10D4	DNA sequence from clone RP1-64K7 on chromosome 20q11.21-11.23 Contains the EIF2S2 gene for eukaryotic translation initiation factor 2 subunit 2 (beta, 38kD), a putative novel gene, the gene for heterogenous nuclear ribonucleoprotein RALY or autoantigen P542, an RPS2 (RPS4) (40S ribosomal protein S2) pseudogene, ESTs, STS, GSSs and two CpG islands	<u>AL031668; 66383-66970</u>	sense
2A6	PAC clone RP3-414A15 from 14q24.3	<u>AC005225; 93459-93782</u>	sense
2D3	DNA sequence from clone RP11-357H24 on chromosome 10	AL451084; 42698-42510; with polyA	antisense
2F8	chromosome 17, clone hRPC.1110_E_20	<u>AC004231; 42223-429716; with polyA</u>	sense
3F11	BAC clone CTD-2314H8	<u>AC079338; 21007-21487</u>	sense
4F7	chromosome 1 clone RP11-109I2	AC091609; 155982-156333	sense
11B8	DNA sequence from clone RP11-735A5 on chromosome 1	AL603888	antisense
12A9	chromosome 18, clone RP11-13N13	AC106037.9	sense

References

- Brenner, S. (1974) The genetics of *Caenorhabditis elegans*. *Genetics* 77:71-94.
- Calabrese, V., Scapagnini, G., Giuffrida Stella, A.M., Bates, T.E. and Clark, J.B. (2001) Mitochondrial involvement in brain function and dysfunction: relevance to aging, neurodegenerative disorders and longevity. *Neurochem Res*, 26: 739-764
- Declercq, W., Logghe, M., Fiers, W. and Contreras, R. (2000) "Cloning and expression of cytokine genes", in the book of *Cytokine Molecular Biology*, Balkwill, F.R. (ed.), IRL Press, Oxford, UK; pp 1-17.
- Ernst, J.F. & Chan, R.K. Characterization of *Saccharomyces cerevisiae* mutants supersensitive to aminoglycoside antibiotics. *J. Bacteriol.* **163**, 8-14 (1985).
- Finkel, T. and Holbrook, N.J. (2000) Oxidants, oxidative stress and the biology of ageing. *Nature* 408:239-47.
- Gershon H. and Gershon D. (2000) The budding yeast, *Saccharomyces cerevisiae*, as a model for aging research: a critical review. *Mech Ageing Dev* 120: 1-22.
- Gietz R. Daniel, and Woods Robin A. (2001) Genetic transformation of yeast. *Biotechniques*. 30:816-20, 822-6, 828 passim.
- Guarente, L. (2001) SIR2 and aging--the exception that proves the rule. *Trends Genet.* 17:391-2.
- Hamilton, M.L., Van Remmen, H., Drake, J.A., Yang, H., Guo, Z.M., Kewitt, K., Walter, C.A. and Richardson, A. (2001) Does oxidative damage to DNA increase with age. *Proc Natl Acad Sci* 98: 10469-10474.
- Ishii, N. (2001) Oxidative stress and aging in *Caenorhabditis elegans*. *Free Radic Res* 33: 857-864.
- Jamieson, D.J., Rivers, S.L. and Stephen, D.W. (1994) Analysis of *Saccharomyces cerevisiae* proteins induced by peroxide and superoxide stress. *Microbiology* 140: 3277-3283.
- Jazwinski, S.M. (1996). Longevity, genes, and aging. *Science* 273: 54-59.
- Jin Y. (1999). Transformation. In *C. elegans – A practical approach*, Hope I.A., ed. (New York, NY, Oxford University Press), pp. 69-96.
- Johnson, F.B., Sinclair, D.A. and Guarente, L. (1999). Molecular biology of aging. *Cell*, 96: 291-302.

- Kennedy, B.K., Austriaco, N.R., and Guarente, L. (1994). Daughter cells of *S. cerevisiae* from old mothers display reduced life span. *J. Cell Bio.* 127, 1985-1993
- 5 - Laun, P., Pichova, A., Madeo, F., Fuchs, J., Ellinger, A., Kohlwein, S., Dawes, I., Frohlich, K.U. and Breitenbach, M. (2001) Aged mother cells of *Saccharomyces cerevisiae* show markers of oxidative stress and aging.
- Martin, G.M., Austad, S.N. and Johnson, T.E. (1996) Genetic analysis of ageing: role of oxidative damage and environmental stresses. *Nat Genet* 13:25-34.
- 10 - Rice R.H. and Etzler M.E. (1975) Chemical modification and hybridization of wheat germ agglutinins. *Biochemistry.* 14:4093-4099.
- Shan, F., Nowell, T.R. Jr. and Taylor, A. (2001) Removal of oxidatively damaged proteins from lens cells by the ubiquitin-proteasome pathway. *Exp Eye Res* 73: 229-238.
- 15 - Smeal T, Claus J, Kennedy B, Cole F, Guarente L. (1996) Loss of transcriptional silencing causes sterility in old mother cells of *S. cerevisiae*. *Cell.* 84:633-42.
- Tanaka, T., Nakamura, H., Nishiyama, A., Hosoi, F., Masutani, H., Wada, H. and Yodoi, J. (2001) Redox regulation by thioredoxin superfamily; protection against oxidative stress and aging. *Free Radic Res* 33, 851-855.
- 20 - Yamamoto, K., Tsuji, T., Matsumoto, I and Osawa, T. (1981) Structural requirements for the binding of oligosaccharides and glycopeptides to immobilized wheat germ agglutinin. *Biochemistry.* 20: 5894-5899.
- Wright C.S., (1984) Structural comparison of the two distinct sugar binding sites in wheat germ agglutinin isolectin II. *J Mol Biol* 178:91-104.

25

Claims

1. Method to screen genes involved in aging and/or in AAD's and/or in oxidative stress, comprising a) mutation or transformation of a yeast cell b) cultivation of said cell c) enrichment of the population for mother cells d) labelling said mother cells with a WGA based label and e) isolation of the highly labelled cells.
2. A method according to claim 1, whereby said WGA based label is FITC-conjugated WGA.
3. A method according to claim 1 or 2, whereby said isolation is a FACS based sorting.
4. A method according to any of the preceding claims, whereby said enrichment is a magnetic-based sorting.
5. A method according to any of the preceding claims, whereby said transformation is carried out with a yeast expression library.
6. A method according to claim 5, whereby said yeast expression library is expressing mammalian DNA or plant DNA.
7. A gene or functional gene fragment isolated with a method according to any of the claims 1-6.
8. A gene or functional gene fragment according to claim 7, comprising SEQ ID N°1, 3, 5, 7, 8, 9, 11, 13, 15, 16, 17, 19, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52 or 53.
9. The use of a gene or functional gene fragment, according to claim 7 or 8, to modulate aging and/or to protect against oxidative stress.
10. The use according to claim 9, whereby said gene comprises SEQ ID N° 11 or 16.
11. A polypeptide, encoded by a functional gene fragment according to claim 7.
12. The use of a polypeptide, encoded by a gene or functional gene fragment according to claim 7, to modulate aging and/or to protect against oxidative stress.
13. The use of a polypeptide, according to claim 11, whereby said gene or functional gene fragment comprises SEQ ID N° 2, 4, 6, 10, 14, 18 or 20.
14. The use of a polypeptide, according to claim 11, whereby said polypeptide comprises SEQ ID N° 12.
15. The use of a polypeptide, according to claim 11, whereby said polypeptide is encoded by SEQ ID N° 16.

1/7

Fig. 1





Fig. 2



3/7

Fig. 3






Fig. 4



4/7

Fig. 5



5/7

Fig. 6





Fig. 7



6/7

Fig. 8






Fig. 9



7/7

Fig. 10



SEQUENCE LISTING

<110> Vlaams Interuniversitair Instituut voor Biotechnol

<120> Method to isolate genes involved in aging

<130> RCO/FAC/V098

<140>

<141>

<150> EP01204600.9

<151> 2001-11-29

<160> 55

<170> PatentIn Ver. 2.1

<210> 1

<211> 1551

<212> DNA

<213> Homo sapiens

<220>

<223> human HN1 cDNA

<220>

<221> CDS

<222> (952)..(1026)

<400> 1

caaaccctact ccaccttact accagacaac cttagccaaa ccatttaccc aaataaagta 60
taggcgatag aaattgaaac ctggcgcaat agatatagta ccgcaaggga aagatgaaaa 120
attataacca agcataatat agcaaggact aaccctata ctttctgcat aatgaattaa 180
ctagaaataa ctttgcaagg agagccaaag ctaagacccc cgaaaccaga cgagctacct 240
aagaacagct aaaagagcac acccgtctat gtagcaaaat agtggaaga tttataggta 300
gaggcgacaa acctaccgag cctggtgata gctggttgtc caagatagaa tcttagttca 360
actttaaatt tgccacaga accctctaaa tccccttgta aatttaactg ttagtccaaa 420
gaggaacagc tntttggaca ctaggaaaaa accttgtaga gagagtaaaa aatttaacac 480
ccatagtagg cctaaaagca gccaccaatt aagaatagcg ttcaagctca acaccacta 540


```

cctaaaaaaaaa tcccaaacaat ataactgaac tcctcacacc caattggacc aatctatcac 600
cctatagaag aactaatggt agtataagta acatgaaaac attctcctcc gcataagcct 660
gcgtcagatt aaaacactga actgacaatt aacagcccaa tatctacaat caaccaacaa 720
gtcattatta ccctcaactgt caacccaaca caggcatgct cataaggaaa gggttaaaaaa 780
agtaaaagga actcggcaaa tcttaccocg cctgtttacc aaaaacatca cctctagcat 840
caccagtatt agaggcaccg cctgccaggt gacacatggt taacggccgc gagtacccta 900
accgtgcaaa ggtagcataa tcacttggtc cttaattagg gacctgtatg a atg gct 957
                                                    Met Ala
                                                    1

cca cga ggg ttc agc tgt ctc tta ctt tta acc agt gaa att gac ctg 1005
Pro Arg Gly Phe Ser Cys Leu Leu Leu Leu Thr Ser Glu Ile Asp Leu
           5                10                15

ccc gtg aag agg cgg gca tga cacagcaaga cgagaagacc ctatggagct 1056
Pro Val Lys Arg Arg Ala
           20                25

ttaatttatt aatgcaaaca gtacctaaca aaccanagg tcctaaacta ccaaacctgc 1116
attaaaaatt tcggttgggg cgacctcgga gcagaacca acctccgagc agtacatgct 1176
aagacttcac cagtcaaagc gaactactat actcaattga tccaataact tgaccaacgg 1236
aacaagttac cctagggata acagcgcaat cctattccta gagtccatat caacaatagg 1296
gtttacgacc tcgatgttgg atcaggacat cccgatggtg cagccgctat taaaggttcg 1356
tttgttcaac gattaaagtc ctacgtgac tgagttcaga ccggagtaat ccaggtcggt 1416
ttctatctac ttcaaattcc tccctgtncg aaaggacaag agaaataagg cctacttcac 1476
aaagcgcctt cccocgtaaa tgatatcatc tcaacttagt attataccca caccaccca 1536
agaacagggt ttggtt 1551

```

<210> 2

<211> 24

<212> PRT

<213> Homo sapiens

<223> human HN1 cDNA

<400> 2

Met Ala Pro Arg Gly Phe Ser Cys Leu Leu Leu Leu Thr Ser Glu Ile
 1 5 10 15
 Asp Leu Pro Val Lys Arg Arg Ala
 20

<210> 3

<211> 884

<212> DNA

<213> Homo sapiens

<220>

<223> human APOA1 cDNA fragment

<220>

<221> CDS

<222> (2)..(829)

<400> 3

a gga ggt ccc cca cgg ccc ttc agg atg aaa gct gcg gtg ctg acc ttg 49
 Gly Gly Pro Pro Arg Pro Phe Arg Met Lys Ala Ala Val Leu Thr Leu
 1 5 10 15

gcc gtg ctc ttc ctg acg ggg agc cag gct cgg cat ttc tgg cag caa 97
 Ala Val Leu Phe Leu Thr Gly Ser Gln Ala Arg His Phe Trp Gln Gln
 20 25 30

gat gaa ccc ccc cag agc ccc tgg gat cga gtg aag gac ctg gcc act 145
 Asp Glu Pro Pro Gln Ser Pro Trp Asp Arg Val Lys Asp Leu Ala Thr
 35 40 45

gtg tac gtg gat gtg ctc aaa gac agc ggc aga gac tat gtg tcc cag 193
 Val Tyr Val Asp Val Leu Lys Asp Ser Gly Arg Asp Tyr Val Ser Gln
 50 55 60

ttt gaa ggc tcc gcc ttg gga aaa cag cta aac cta aag ctc ctt gac 241
 Phe Glu Gly Ser Ala Leu Gly Lys Gln Leu Asn Leu Lys Leu Leu Asp
 65 70 75 80

aac tgg gac agc gtg acc tcc acc ttc agc aag ctg cgc gaa cag ctc 289
 Asn Trp Asp Ser Val Thr Ser Thr Phe Ser Lys Leu Arg Glu Gln Leu
 85 90 95

ggc cct gtg acc cag gag ttc tgg gat aac ctg gaa aag gag aca gag 337

Gly	Pro	Val	Thr	Gln	Glu	Phe	Trp	Asp	Asn	Leu	Glu	Lys	Glu	Thr	Glu		
			100					105					110				
ggc	ctg	agg	cag	gag	atg	agc	aag	gat	ctg	gag	gag	gtg	aag	gcc	aag	385	
Gly	Leu	Arg	Gln	Glu	Met	Ser	Lys	Asp	Leu	Glu	Glu	Val	Lys	Ala	Lys		
		115					120					125					
gtg	cag	ccc	tac	ctg	gac	gac	ttc	cag	aag	aag	tgg	cag	gag	gag	atg	433	
Val	Gln	Pro	Tyr	Leu	Asp	Asp	Phe	Gln	Lys	Lys	Trp	Gln	Glu	Glu	Met		
		130					135				140						
gag	ctc	tac	cgc	cag	aag	gtg	gag	ccg	ctg	cgc	gca	gag	ctc	caa	gag	481	
Glu	Leu	Tyr	Arg	Gln	Lys	Val	Glu	Pro	Leu	Arg	Ala	Glu	Leu	Gln	Glu		
145					150				155					160			
ggc	gcg	cgc	cag	aag	ctg	cac	gag	ctg	caa	gag	aag	ctg	agc	cca	ctg	529	
Gly	Ala	Arg	Gln	Lys	Leu	His	Glu	Leu	Gln	Glu	Lys	Leu	Ser	Pro	Leu		
			165					170					175				
ggc	gag	gag	atg	cgc	gac	cgc	gcg	cgc	gcc	cat	gtg	gac	gcg	ctg	cgc	577	
Gly	Glu	Glu	Met	Arg	Asp	Arg	Ala	Arg	Ala	His	Val	Asp	Ala	Leu	Arg		
			180					185					190				
acg	cat	ctg	gcc	ccc	tac	agc	gac	gag	ctg	cgc	cag	cgc	ttg	gcc	gcg	625	
Thr	His	Leu	Ala	Pro	Tyr	Ser	Asp	Glu	Leu	Arg	Gln	Arg	Leu	Ala	Ala		
		195						200				205					
cgc	ctt	gag	gct	ctc	aag	gag	aac	ggc	ggc	gcc	aga	ctg	gcc	gag	tac	673	
Arg	Leu	Glu	Ala	Leu	Lys	Glu	Asn	Gly	Gly	Ala	Arg	Leu	Ala	Glu	Tyr		
		210					215				220						
cac	gcc	aag	gcc	acc	gag	cat	ctg	agc	acg	ctc	agc	gag	aag	gcc	aag	721	
His	Ala	Lys	Ala	Thr	Glu	His	Leu	Ser	Thr	Leu	Ser	Glu	Lys	Ala	Lys		
225					230					235				240			
ccc	gcg	ctc	gag	gac	ctc	cgc	caa	ggc	ctg	ctg	ccc	gtg	ctg	gag	agc	769	
Pro	Ala	Leu	Glu	Asp	Leu	Arg	Gln	Gly	Leu	Leu	Pro	Val	Leu	Glu	Ser		
				245				250					255				
ttc	aag	gtc	agc	ttc	ctg	agc	gct	ctc	gag	gag	tac	act	aag	aag	ctc	817	
Phe	Lys	Val	Ser	Phe	Leu	Ser	Ala	Leu	Glu	Glu	Tyr	Thr	Lys	Lys	Leu		
			260					265					270				
aac	acc	cag	tga	ggcgcccgcc	gccgcccccc	ttcccggtgc	tcagaataaa									869	
Asn	Thr	Gln															
		275															
cgtttcctaaa	gtggg															884	

<210> 4
 <211> 275
 <212> PRT
 <213> Homo sapiens
 <223> human APOA1 cDNA fragment

<400> 4

```

Gly Gly Pro Pro Arg Pro Phe Arg Met Lys Ala Ala Val Leu Thr Leu
 1           5           10           15
Ala Val Leu Phe Leu Thr Gly Ser Gln Ala Arg His Phe Trp Gln Gln
          20           25           30
Asp Glu Pro Pro Gln Ser Pro Trp Asp Arg Val Lys Asp Leu Ala Thr
          35           40           45
Val Tyr Val Asp Val Leu Lys Asp Ser Gly Arg Asp Tyr Val Ser Gln
          50           55           60
Phe Glu Gly Ser Ala Leu Gly Lys Gln Leu Asn Leu Lys Leu Leu Asp
          65           70           75           80
Asn Trp Asp Ser Val Thr Ser Thr Phe Ser Lys Leu Arg Glu Gln Leu
          85           90           95
Gly Pro Val Thr Gln Glu Phe Trp Asp Asn Leu Glu Lys Glu Thr Glu
          100          105          110
Gly Leu Arg Gln Glu Met Ser Lys Asp Leu Glu Glu Val Lys Ala Lys
          115          120          125
Val Gln Pro Tyr Leu Asp Asp Phe Gln Lys Lys Trp Gln Glu Glu Met
          130          135          140
Glu Leu Tyr Arg Gln Lys Val Glu Pro Leu Arg Ala Glu Leu Gln Glu
          145          150          155          160
Gly Ala Arg Gln Lys Leu His Glu Leu Gln Glu Lys Leu Ser Pro Leu
          165          170          175
Gly Glu Glu Met Arg Asp Arg Ala Arg Ala His Val Asp Ala Leu Arg
          180          185          190
Thr His Leu Ala Pro Tyr Ser Asp Glu Leu Arg Gln Arg Leu Ala Ala
          195          200          205
Arg Leu Glu Ala Leu Lys Glu Asn Gly Gly Ala Arg Leu Ala Glu Tyr
          210          215          220
His Ala Lys Ala Thr Glu His Leu Ser Thr Leu Ser Glu Lys Ala Lys
          225          230          235          240
Pro Ala Leu Glu Asp Leu Arg Gln Gly Leu Leu Pro Val Leu Glu Ser
          245          250          255
Phe Lys Val Ser Phe Leu Ser Ala Leu Glu Glu Tyr Thr Lys Lys Leu
          260          265          270
Asn Thr Gln
          275

```

<210> 5

<211> 771

<212> DNA

<213> Homo sapiens

<220>

<223> human ribosomal protein P O cDNA fragment

<220>

<221> CDS

<222> (10)..(705)

<400> 5

cggatatga ggc agc agt ttc tcc aag gag gac ctc act gag atc agg gac 51
 Gly Ser Ser Phe Ser Lys Glu Asp Leu Thr Glu Ile Arg Asp
 1 5 10

atg ttg ctg gcc aat aag gtg cca gct gct gcc cgt gct ggt gcc att 99
 Met Leu Leu Ala Asn Lys Val Pro Ala Ala Ala Arg Ala Gly Ala Ile
 15 20 25 30

gcc cca tgt gaa gtc act gtg cca gcc cag aac act ggt ctc ggg ccc 147
 Ala Pro Cys Glu Val Thr Val Pro Ala Gln Asn Thr Gly Leu Gly Pro
 35 40 45

gag aag acc tcc ttt ttc cag gct tta ggt atc acc act aaa atc tcc 195
 Glu Lys Thr Ser Phe Phe Gln Ala Leu Gly Ile Thr Thr Lys Ile Ser
 50 55 60

agg ggc acc att gaa atc ctg agt gat gtg cag ctg atc aag act gga 243
 Arg Gly Thr Ile Glu Ile Leu Ser Asp Val Gln Leu Ile Lys Thr Gly
 65 70 75

gac aaa gtg gga gcc agc gaa gcc acg ctg ctg aac atg ctc aac atc 291
 Asp Lys Val Gly Ala Ser Glu Ala Thr Leu Leu Asn Met Leu Asn Ile
 80 85 90

tcc ccc ttc tcc ttt ggg ctg gtc acc cag cag gtg ttc gac aat ggc 339
 Ser Pro Phe Ser Phe Gly Leu Val Thr Gln Gln Val Phe Asp Asn Gly
 95 100 105 110

agc atc tac aac cct gaa gtg ctt gat atc aca gag gaa act ctg cat 387
 Ser Ile Tyr Asn Pro Glu Val Leu Asp Ile Thr Glu Glu Thr Leu His
 115 120 125

tct cgc ttc ctg gag ggt gtc cgc aat gtt gcc agt gtc tgt ctg cag 435
 Ser Arg Phe Leu Glu Gly Val Arg Asn Val Ala Ser Val Cys Leu Gln
 130 135 140

```

att ggc tac cca act gtt gca tca gta ccc cat tct atc atc aac ggg      483
Ile Gly Tyr Pro Thr Val Ala Ser Val Pro His Ser Ile Ile Asn Gly
      145                      150                      155

tac aaa cga gtc ctg gcc ttg tct gtg gag acg gat tac acc ttc cca      531
Tyr Lys Arg Val Leu Ala Leu Ser Val Glu Thr Asp Tyr Thr Phe Pro
      160                      165                      170

ctt gct gaa aag gtc aag gcc ttc ttg gct gat cca tct gcc ttt gtg      579
Leu Ala Glu Lys Val Lys Ala Phe Leu Ala Asp Pro Ser Ala Phe Val
      175                      180                      185                      190

gct gct gcc cct gtg gct gct gcc acc aca gct gct cct gct gct gct      627
Ala Ala Ala Pro Val Ala Ala Ala Thr Thr Ala Ala Pro Ala Ala Ala
      195                      200                      205

gca gcc cca gct aag gtt gaa gcc aag gaa gag tcg gag gag tcg gac      675
Ala Ala Pro Ala Lys Val Glu Ala Lys Glu Glu Ser Glu Glu Ser Asp
      210                      215                      220

gag gat atg gga ttt ggt ctc ttt gac taa tcaccaaaaa gcaaccaact      725
Glu Asp Met Gly Phe Gly Leu Phe Asp
      225                      230

tagccagttt tatttgcaaa acaaggaaat aaaggcttac ttcttt      771

```

<210> 6

<211> 231

<212> PRT

<213> Homo sapiens

<223> human ribosomal protein P O cDNA fragment

<400> 6

```

Gly Ser Ser Phe Ser Lys Glu Asp Leu Thr Glu Ile Arg Asp Met Leu
  1                      5                      10                      15
Leu Ala Asn Lys Val Pro Ala Ala Ala Arg Ala Gly Ala Ile Ala Pro
      20                      25                      30
Cys Glu Val Thr Val Pro Ala Gln Asn Thr Gly Leu Gly Pro Glu Lys
      35                      40                      45
Thr Ser Phe Phe Gln Ala Leu Gly Ile Thr Thr Lys Ile Ser Arg Gly
      50                      55                      60
Thr Ile Glu Ile Leu Ser Asp Val Gln Leu Ile Lys Thr Gly Asp Lys
      65                      70                      75                      80
Val Gly Ala Ser Glu Ala Thr Leu Leu Asn Met Leu Asn Ile Ser Pro
      85                      90                      95
Phe Ser Phe Gly Leu Val Thr Gln Gln Val Phe Asp Asn Gly Ser Ile

```

100	105	110
Tyr Asn Pro Glu Val Leu Asp Ile Thr Glu Glu Thr Leu His Ser Arg		
115	120	125
Phe Leu Glu Gly Val Arg Asn Val Ala Ser Val Cys Leu Gln Ile Gly		
130	135	140
Tyr Pro Thr Val Ala Ser Val Pro His Ser Ile Ile Asn Gly Tyr Lys		
145	150	155
Arg Val Leu Ala Leu Ser Val Glu Thr Asp Tyr Thr Phe Pro Leu Ala		
165	170	175
Glu Lys Val Lys Ala Phe Leu Ala Asp Pro Ser Ala Phe Val Ala Ala		
180	185	190
Ala Pro Val Ala Ala Ala Thr Thr Ala Ala Pro Ala Ala Ala Ala Ala		
195	200	205
Pro Ala Lys Val Glu Ala Lys Glu Glu Ser Glu Glu Ser Asp Glu Asp		
210	215	220
Met Gly Phe Gly Leu Phe Asp		
225	230	

<210> 7

<211> 2769

<212> DNA

<213> Homo sapiens

<220>

<223> human glutamyl tRNA synthase antisense cDNA

<400> 7

```

agaatttagc tgttctttat tgacatggaa tttgggggtg cgagggtagc cacaccctcc 60
aggaggatga ggtaggttca gtgcttccag ctcacacctt tcctgggtct tccttcagtg 120
tgacagttcg gttaaagaca agctttccct gatggctgtc tggatccacg gagaaatata 180
caagacgctc aaactggaac ttgtcgaagg gttttgccag ggccacagag cagtccacta 240
atgctgcata caccacgtgt agtgatgcca ggttcaggct acttaaaaaat ccaccaggca 300
cctcagtagg atcttcaggg ttcttgtgct ggaatagtcg ctcatagagg cgaacctcac 360
acatcaaagg ctgtgacacc cagtgaataa aggccttttg cttctctcca gcatctgccc 420
gtctgcaggt cacctccaga ctctctacac aacctctggg gcccttgaca acatgctgca 480
gctcaatgac gtagcctgta tgccctcaggc ccacaggctg gcccgaagcc aggcgcttaa 540
atcctggctc tggctcctcc ttgaagtcag tcctctcaat gaagacaatg ggtgcaaagg 600
gaacctgatg gaagcctttg gtctcatcag ctgggaagtt gggcacctgg atgtccaagg 660
acttggcagc aggaaagttg gtgatgatga cccgtagtga ctccagcaca gccatggctc 720
gtggggctgt gtcattcagc acatcacgca cacaggcttc tagaagatgt ggctccattg 780
tggtttgtgc cacagtcact cccaccggg cacagaagtt gttgatggcc tcaggtggga 840
agccccgccg tcgcagggcc gtgagtgtaa agagccgtgg gtcaccccag tcccgcacag 900
caccagttgc tacaagctgg aggatcttcc tcttagagac aacagcatag tgcaggttga 960
ggcggccata ctcccactgc acagggcaat agacgtncag tgcattgcaa agccagaagt 1020
aggaagagcg tcgggcctgg aattccttgg tgcanagtga gtgagtgatg tgctcgatgg 1080
agtcacagag gcagtgtgtg tagtcgtagg tgggtagatg gcaccatttg tcccctgtgc 1140

```

```

ggtggtgtgg tgtatacttg actcgatagg ctacaggggc catcttgcca tcctccatca 1200
ccagcttcat ccgtagtgtg gcctcgccct ctgaaaactt gcccttgccg attgcctcaa 1260
agagcagcag tgactcctcc atgggacggt ctctccaggg tgaaggcaga gtattatggc 1320
ctttgagctc ctctcctcgc tgggtggcaca cataagccag acccctgcgg atgagctcca 1380
cagcccacgc atatagctgg tcaaaatagt cagacgcata tgtgactttg taagggtgtgt 1440
agcctagcca ggctaccatg tcacagatgg ccgtgaagaa ctttgcttcc tccttctcag 1500
ggttggtgtc atcaaaaacgc agaaaacaga tgccattggt gcccttggca tagccaaagt 1560
tgaaattgat ggctttggca tgtccaatat gcaggattcc attgggttct ggcggaacc 1620
gggtacgtac ctgcccacca ggttgcttcc gaacatcctg ctcaccggta caccaggggt 1680
tggaataacc acactaggca aagaacttgc gtcaaaatca ggactgaaat acattaatgt 1740
gggtgattta gctcgagaag tctgatcatc ggatatcatg gagtctggca agacggcttc 1800
tcccaagagc atgccgaaag atgcacagat gatggcaca atcctgaagg atatggggat 1860
tacagaatat gagccaagag ttataaatca gatgttggag tttgccttcc gatatgtgac 1920
cacaattcta gatgatgcaa aaatttatcc aagccatgct aagaaagcta ctgttgatgc 1980
agatgatgtg cgattggcaa tccagtgccg cgctgatcag tcttttacct ctctccccc 2040
aagagatttt ttattagata ttgcaaggca aagaaatcaa acccctttgc cattgatcaa 2100
gccatattca ggtcctaggt tgccacctga tagatactgc ttaacagctc caaactatag 2160
gctgaaatct ttacagaaaa aggcataaac ttctgcggga agaataacag tcccgcggtt 2220
aagtgttggg tcagttacta gcagaccaag tactccaca ctaggcacac caaccccaca 2280
gaccatgtct gtttcaacta aagtagggac tcccatgtcc ctcacaggtc aaagggtttac 2340
agtacagatg cctacttctc agtctccagc tgtaaaagct tcaattcctg caacctcagc 2400
agttcagaat gttctgatta atccatcatt aatcgggtcc aaaaacattc ttattaccac 2460
taatatgatg tcatcacaaa atactgccaa tgaatcatca aatgcattga aaagaaaacg 2520
tgaagatgat gatgatgacg atgatgatga tgatgactat gataatctgt aatctagcct 2580
tgctgaatgt aacatgtata cttggtcttg aattcattgt actnatatta aacatgcatg 2640
ctggatgttt tcaagttgtg ttttagaaaa ctaataataa tgagtaaaca cagttaccat 2700
acttttcaat tgaaatgaag gtttttcatc agccttaaaa gtgtaagaaa aataaagttg 2760
tcattcatt
2769

```

<210> 8

<211> 2233

<212> DNA

<213> Homo sapiens

<220>

<223> human G rich sequence factor antisense cDNA

<400> 8

```

tgcacataaa gttagtttat taatgactat attttgaagc cagccatttt gtccaatatt 60
taaataacaa gctgtttaat attaaagcag aaagtactgc cacattgtga cagaagtaca 120
gctttatcca taaacccttc acacaattat acattaaatg ctatttttat ttaagcaagg 180
caccctact tgttctaaaa tatgggatgt actactccat ttaaaaagca aatgaggaga 240
ctgatttttt tcctatctag agctggttta aattcaagtg aagccattaa ttttcttacc 300
agtctggact gttctgacat gtcacttcac agttttgaga ctaacaaaca cccttaggtc 360
taccocaaac caaaaaaagc cagggaggga cagtttanac aacttttaag ntaagactag 420
aatgcccccn taagtagata ggcaccaccg ttatagcctc ttagcacaag gcttagaaaag 480
gcaaaacatt tctaaagaca taaaaccgcc aaattgtcat ttacactaag ctgtggtatt 540

```



```

ttttgttttg atttttataaa attcctatct ctaagtaatc aaatacagca aaatttataaa 600
ctattctgcc aaaccatacc agttcaaagg tctgaaaccg ttctttgctt tggatgaatgt 660
ttgggtataaa taaatcactg ttctactcca tgagaagttt tacatcagag cactcaattc 720
acaggcaaaa aaaaaaagtt nctttttataa tgaaagcggt taaagaacac tgctgaataa 780
aaatatgtgg tctttacaat gagtaacata tcatctatga aaacaaacca tttaatacata 840
taaaacaaag accatatttt taaatcagat cctggacagt ttaaataaaa attttaattt 900
gtgggtttatt tcataatccc aaactggatc ttctatttct tcttctaaac agtccactat 960
gaaacttttc ttagaagcaa ttaacccaaa acacccaaat catttcccta gcagttgctg 1020
ctaataaact ggaactgtat atcccaagtc caagaaagat gtgcaaataa aatgcttctt 1080
gcttcacctt tattatctgg agcccctaga gtcttatttt ccttttggac atgaattcag 1140
gaacagttca atatacctat gatgaacgtg ggaccgatcc ttgagcatcg ctgcaacagc 1200
atcctcatgg gtctcaaagt gcacatcagc ttctccagtg gccttcccac tggagctgta 1260
ttccatgggt attctaacag gcttgagtggt agcaaaaaag ttataaatgt cttgggcatt 1320
ggcttggaag ggtaatcctc tcatgtggac aaaatgcaga gaagacgtag ttccaaaatc 1380
agcagcctct ggaagctttt ctggcacctc cttaggcaat tctatttctt tctcactttc 1440
aaaagctgtc atgggttgaa tatcctcatt tacttcatgt tcttcaaaga ccatttcttg 1500
ctcagttata tacttagcag taggaaaaga tgcgattttc ttcccttat aagaaccgac 1560
atgtgttcga acttcattcc ttctgcttgg aaatatctcg atgtatcgat taccaatttc 1620
ttccctgtgt ttcaacaggg cttggttggc catttctggt tcttcaaatt gcacataggc 1680
ttccctgtgt ttctgcctcc ctctatagtc catcacaata gtaatgtcaa ctatattcag 1740
tcctgcaaaag aagtctacaa tgtctttctc attgcaacta taaggaagtc ctctcaaacg 1800
aaccacacca tcatttacca caggcgaaga ttgacctgc aagctcttca ttaaggcatc 1860
cacatcttca ttgtttatct catatacttc cacataccgc tggcccatgt acatacgggtg 1920
cttctctaag gctttntgca catcctgctc tgactccatt tcaattaagg catcaccctt 1980
tcgtttccca tctctgttta ggagaaaatg tattccattc tcaccgttgc ggattctgca 2040
gtctgaaaaa aagttaagca catcttccat agtgcagac cagggcagtc cttgagctcg 2100
aatgagaaag acatcatcca ctctctctc taacttgac ggggccaatt catactcagg 2160
gggtggtgga aggtcttcca ggtaagtagt ttgggactcc tggctgtagc tgcgcgtcgg 2220
gacggcggcc gcc 2233

```

<210> 9

<211> 593

<212> DNA

<213> Homo sapiens

<220>

<223> human ALDH1 cDNA fragment

<220>

<221> CDS

<222> (2)..(55)

<400> 9

t gaa tat aca gag gtc aaa aca gtc aca gtg aaa atc tct cag aag aac 49

Glu Tyr Thr Glu Val Lys Thr Val Thr Val Lys Ile Ser Gln Lys Asn

1

5

10

15

tca taa agaaaataca agagtggaga gaagctcttc aatagctaag catctcctta 105
 Ser

cagtcactaa tatagtagat tttaaagaca aaatTTTTTct tttcttgatt tttttaaaca 165

taagctaaat catattagta ttaatactac ccatagaaaa cttgacatgt agcttcttct 225

gaaagaatta tttgccttct gaaatgtgac cccaagtcc tatcctaaat aaaaaagac 285

aaattcggat gtatgatctc tctagctttg tcatagttat gtgattttcc tttgtagcta 345

cttttgcagg ataataattt tatagaaaag gaacagttgc atttagcttc tttcccttag 405

tgactcttga agtacttaac atacacgtta actgcagagt aaattgctct gttcccagta 465

gttataaagt ctttgactg ttttgaaaag tttcctagga tgtcatgtct gottgtcaaa 525

agaaataatc cctgtaatat ttagctgtaa actgaatata aagcttaata aaaacaacct 585

tgcatgat 593

<210> 10
 <211> 17
 <212> PRT
 <213> Homo sapiens
 <223> human ALDH1 cDNA fragment

<400> 10
 Glu Tyr Thr Glu Val Lys Thr Val Thr Val Lys Ile Ser Gln Lys Asn
 1 5 10 15
 Ser

<210> 11
 <211> 418
 <212> DNA
 <213> Homo sapiens

<220>
 <223> human ferritin cDNA fragment

<220>
 <221> CDS
 <222> (2)..(418)

<400> 11

```

c ggt ccc gcg ggt ctg tct ctt gct tca aca gtg ttt gga cgg aac aga 49
  Gly Pro Ala Gly Leu Ser Leu Ala Ser Thr Val Phe Gly Arg Asn Arg
    1             5             10             15

tcc ggg gac tct ctt cca gcc tcc gac cgc cct ccg att tcc tct ccg 97
  Ser Gly Asp Ser Leu Pro Ala Ser Asp Arg Pro Pro Ile Ser Ser Pro
           20             25             30

ctt gca acc tcc ggg acc atc ttc tcg gcc atc tcc tgc ttc tgg gac 145
  Leu Ala Thr Ser Gly Thr Ile Phe Ser Ala Ile Ser Cys Phe Trp Asp
           35             40             45

ctg cca gca ccg ttt ttg tgg tta gct cct tct tgc caa cca acc atg 193
  Leu Pro Ala Pro Phe Leu Trp Leu Ala Pro Ser Cys Gln Pro Thr Met
           50             55             60

agc tcc cag att cgt cag aat tat tcc acc gac gtg gag gca gcc gtc 241
  Ser Ser Gln Ile Arg Gln Asn Tyr Ser Thr Asp Val Glu Ala Ala Val
        65             70             75             80

aac agc ctg gtc aat ttg tac ctg cag gcc tcc tac acc tac ctc tct 289
  Asn Ser Leu Val Asn Leu Tyr Leu Gln Ala Ser Tyr Thr Tyr Leu Ser
           85             90             95

ctg ggc ttc tat ttc gac cgc gat gat gtg gct ctg gaa ggc gtg agc 337
  Leu Gly Phe Tyr Phe Asp Arg Asp Asp Val Ala Leu Glu Gly Val Ser
           100             105             110

cac ttc ttc cgc gaa ttg gcc gag gag aag cgc gag ggc tac gag cgt 385
  His Phe Phe Arg Glu Leu Ala Glu Glu Lys Arg Glu Gly Tyr Glu Arg
           115             120             125

ctc ctg aag atg caa aac cag cgt ggc ggc cgc 418
  Leu Leu Lys Met Gln Asn Gln Arg Gly Gly Arg
        130             135

```

<210> 12

<211> 139

<212> PRT

<213> Homo sapiens

<223> human ferritin cDNA fragment

<400> 12

```

Gly Pro Ala Gly Leu Ser Leu Ala Ser Thr Val Phe Gly Arg Asn Arg
  1             5             10             15

```

```

Ser Gly Asp Ser Leu Pro Ala Ser Asp Arg Pro Pro Ile Ser Ser Pro

```

20 25 30
 Leu Ala Thr Ser Gly Thr Ile Phe Ser Ala Ile Ser Cys Phe Trp Asp
 35 40 45
 Leu Pro Ala Pro Phe Leu Trp Leu Ala Pro Ser Cys Gln Pro Thr Met
 50 55 60
 Ser Ser Gln Ile Arg Gln Asn Tyr Ser Thr Asp Val Glu Ala Ala Val
 65 70 75 80
 Asn Ser Leu Val Asn Leu Tyr Leu Gln Ala Ser Tyr Thr Tyr Leu Ser
 85 90 95
 Leu Gly Phe Tyr Phe Asp Arg Asp Asp Val Ala Leu Glu Gly Val Ser
 100 105 110
 His Phe Phe Arg Glu Leu Ala Glu Glu Lys Arg Glu Gly Tyr Glu Arg
 115 120 125
 Leu Leu Lys Met Gln Asn Gln Arg Gly Gly Arg
 130 135

<210> 13

<211> 929

<212> DNA

<213> Homo sapiens

<220>

<223> human ribosomal protein S2 cDNA

<220>

<221> CDS

<222> (11)..(892)

<400> 13

aaaacaccaa atg gcg gat gac gcc ggt gca gcg ggg ggg ccc ggg ggc 49
 Met Ala Asp Asp Ala Gly Ala Ala Gly Gly Pro Gly Gly
 1 5 10
 cct ggt ggc cct ggg atg ggg aac cgc ggt ggc ttc cgc gga ggt ttc 97
 Pro Gly Gly Pro Gly Met Gly Asn Arg Gly Gly Phe Arg Gly Gly Phe
 15 20 25
 ggc agt ggc atc cgg ggc cgg ggt cgc ggc cgt gga cgg ggc cgg ggc 145
 Gly Ser Gly Ile Arg Gly Arg Gly Arg Gly Arg Gly Arg Gly

30	35	40	45	
cga ggc cgc gga gct cgc gga ggc aag gcc gag gat aag gag tgg atg				193
Arg Gly Arg Gly Ala Arg Gly Gly Lys Ala Glu Asp Lys Glu Trp Met				
	50	55	60	
ccc gtc acc aag ttg ggc cgc ttg gtc aag gac atg aag atc aag tcc				241
Pro Val Thr Lys Leu Gly Arg Leu Val Lys Asp Met Lys Ile Lys Ser				
	65	70	75	
ctg gag gag atc tat ctc ttc tcc ctg ccc att aag gaa tca gag atc				289
Leu Glu Glu Ile Tyr Leu Phe Ser Leu Pro Ile Lys Glu Ser Glu Ile				
	80	85	90	
att gat ttc ttc ctg ggg gcc tct ctc aag gat gag gtt ttg aag att				337
Ile Asp Phe Phe Leu Gly Ala Ser Leu Lys Asp Glu Val Leu Lys Ile				
	95	100	105	
atg cca gtg cag aag cag acc cgt gcc ggc cag cgc acc agg ttc aag				385
Met Pro Val Gln Lys Gln Thr Arg Ala Gly Gln Arg Thr Arg Phe Lys				
	110	115	120	125
gca ttt gtt gct atc ggg gac tac aat ggc cac gtc ggt ctg ggt gtt				433
Ala Phe Val Ala Ile Gly Asp Tyr Asn Gly His Val Gly Leu Gly Val				
	130	135	140	
aag tgc tcc aag gag gtg gcc acc gcc atc cgt ggg gcc atc atc ctg				481
Lys Cys Ser Lys Glu Val Ala Thr Ala Ile Arg Gly Ala Ile Ile Leu				
	145	150	155	
gcc aag ctc tcc atc gtc ccc gtg cgc aga ggc tac tgg ggg aac aag				529
Ala Lys Leu Ser Ile Val Pro Val Arg Arg Gly Tyr Trp Gly Asn Lys				
	160	165	170	
atc ggc aag ccc cac act gtc cct tgc aag gtg aca ggc cgc tgc ggc				577
Ile Gly Lys Pro His Thr Val Pro Cys Lys Val Thr Gly Arg Cys Gly				
	175	180	185	
tct gtg ctg gta cgc ctc atc cct gca ccc agg ggc act ggc atc gtc				625
Ser Val Leu Val Arg Leu Ile Pro Ala Pro Arg Gly Thr Gly Ile Val				
	190	195	200	205
tcc gca cct gtg cct aag aag ctg ctc atg atg gct ggt atc gat gac				673
Ser Ala Pro Val Pro Lys Lys Leu Leu Met Met Ala Gly Ile Asp Asp				
	210	215	220	
tgc tac acc tca gcc cgg ggc tgc act gcc acc ctg ggc aac ttc gcc				721
Cys Tyr Thr Ser Ala Arg Gly Cys Thr Ala Thr Leu Gly Asn Phe Ala				

```

<400> 14
Met Ala Asp Asp Ala Gly Ala Ala Gly Gly Pro Gly Gly Pro Gly Gly
  1          5          10          15
Pro Gly Met Gly Asn Arg Gly Gly Phe Arg Gly Gly Phe Gly Ser Gly
      20          25          30
Ile Arg Gly Arg Gly Arg Gly Arg Gly Arg Gly Arg Gly Arg Gly Arg
      35          40          45
Gly Ala Arg Gly Gly Lys Ala Glu Asp Lys Glu Trp Met Pro Val Thr
      50          55          60
Lys Leu Gly Arg Leu Val Lys Asp Met Lys Ile Lys Ser Leu Glu Glu
  65          70          75          80
Ile Tyr Leu Phe Ser Leu Pro Ile Lys Glu Ser Glu Ile Ile Asp Phe
      85          90          95
Phe Leu Gly Ala Ser Leu Lys Asp Glu Val Leu Lys Ile Met Pro Val
      100          105          110
Gln Lys Gln Thr Arg Ala Gly Gln Arg Thr Arg Phe Lys Ala Phe Val
      115          120          125
Ala Ile Gly Asp Tyr Asn Gly His Val Gly Leu Gly Val Lys Cys Ser
      130          135          140
Lys Glu Val Ala Thr Ala Ile Arg Gly Ala Ile Ile Leu Ala Lys Leu
  145          150          155          160

```

```

Ser Ile Val Pro Val Arg Arg Gly Tyr Trp Gly Asn Lys Ile Gly Lys
      165                      170                      175
Pro His Thr Val Pro Cys Lys Val Thr Gly Arg Cys Gly Ser Val Leu
      180                      185                      190
Val Arg Leu Ile Pro Ala Pro Arg Gly Thr Gly Ile Val Ser Ala Pro
      195                      200                      205
Val Pro Lys Lys Leu Leu Met Met Ala Gly Ile Asp Asp Cys Tyr Thr
      210                      215                      220
Ser Ala Arg Gly Cys Thr Ala Thr Leu Gly Asn Phe Ala Lys Ala Thr
      225                      230                      235                      240
Phe Asp Ala Ile Ser Lys Thr Tyr Ser Tyr Leu Thr Pro Asp Leu Trp
      245                      250                      255
Lys Glu Thr Val Phe Thr Lys Ser Pro Tyr Gln Glu Phe Thr Asp His
      260                      265                      270
Leu Val Lys Thr His Thr Arg Val Ser Val Gln Arg Thr Gln Ala Pro
      275                      280                      285
Ala Val Ala Thr Thr
      290

```

<210> 15

<211> 525

<212> DNA

<213> Homo sapiens

<220>

<223> human H2A histone antisense cDNA

<400> 15

```

ccgctttttt tttttttttt aaaaaaaaaa caatccttta ataaaattag tccatctaaa 60
actccccaat gcctaagggt ctagtcgtgg aagggttagc tgcagaattc cagttcagaa 120
gccaacggag gctcagtcga gacagggatt aaccgacttg tgctggtatc taggtgcttg 180
gattgccgag ttgagtttgc tggaaggga atggggcggc gtcggggggc cgcacaggcc 240
tggtctgccg tcggcttcgc cgaaaacgac tcttgctgtc cacatagcca gccgtgaagc 300
caaactgcag gcctctgccc tcccctaaat gtcctcctag gaggcagtca ctcgggagga 360
agatgtgcct gttaccaagt gcttcgtccc ggccccagcg cagacctatg aatgaagatg 420
gagggagagc tgatgtgaaa ggccctgggtc ccggccgcag aaggggagcc agaccgaaat 480
cggcgatcgc gtctctccca ccaaagcctc ccggccagcg gccgc 525

```

<210> 16

<211> 1381

<212> DNA

<213> Homo sapiens

<400> 16

```

catctggctg taaggtttgc ccttgaacta aaaatgttgt ttggggcaag ggccagaaat 60

```

```

gtggagacat ggtttttgtt acgcattcctt gtattatatg tgactaaatt tacaaacaag 120
atacatgtgt aattaaagac ccttatggaa ctggaagacg tcttgtagtg ctacattggg 180
tgaaaccggt ggtccattttt tgtctgtttc tatgaagata aaataattgg gggccatcta 240
gaaatagaaa ggcagtggga agacagattc tacggcactg ctttcattta attgggcttt 300
aggcactcca ttcgaatgca gaacctcacc tctagttgag accaagaatt ggcaaatttg 360
catgagctcc tggaaagagt tgctgacttt gtatctaaga cctgccaggg aataccaaga 420
gttgtttcta cagactttttt tttttttttt tgtatgggag aagatactgt ggcaaccagg 480
aaggaatgga aaaaaaattc ttttctctac agcaaattaa tgtgaggaag ctctccaat 540
cctctggcta tttaagggtc aaaatcaagt gcctagggaa aattccaatg gatgattttc 600
tgggagctat cttgtctacc ttgagggtcc tgaacaatga attccatta atgagcagtc 660
ttcagtatta aaaccaactgt cttgtcacct ctttttgcac tactgtcttc cgtggatgtt 720
tcagttacaa ctgtaatgtt atttatagaa caacattaat ccattaaagc taacctattt 780
ttcaatattt atgataatct atgtacatat attgtctgtc catatgtatt tgtaaatagg 840
ttgtatataa tgtcagggtt gggctcttggg ttcaagtgtg tatattcctg taagtttctt 900
aactgcattt tgatgaattc acattatgta actataagaa ttgtcccaa agtacctgta 960
cagaaaattg aatattgaaa aattgacaaa ttgtgtacaa aactaaaaa aaacttggtt 1020
aaattgtatt tgcaataaac aacatcaa attttcatga aatcttggtg caaattcaga 1080
tctcttattt aaaatttaaa taaggaatac attttcaaaa tgcagtaatc aaaatgtgat 1140
ctagtgtaat gaaataaaat gtgatctagt gtaatggaag acctttgaga acctgggtgt 1200
attaactttg tgtatatagt gtaaatatcc ccactgtact gttagaggcc aacaattcta 1260
gtatggcttg ttggcaaaga gtgctacacc gtttcaatga aacaatgtat gtttgtttta 1320
actgaactaa aataaataca tgcttaatcc cgaaaaaaaa aaaaacaaaa aaaaaaaaaa 1380
a                                                                                   1381

```

<210> 17

<211> 2721

<212> DNA

<213> Homo sapiens

<220>

<223> clone 11A12 - hypothetical protein DKFZp564K0822

<220>

<221> CDS

<222> (75)..(581)

<400> 17

```

gggctgctct tggagtgcac agaagccaaa aagcattgct ggtatttcga aggactctat 60

ccaacctatt atat atg ccg ctc cta cga gga ctg ctg tgg ctc cag gtg 110
          Met Pro Leu Leu Arg Gly Leu Leu Trp Leu Gln Val
                1                      5                      10

ctg tgt gcg ggc cct ctc cat aca gag gct gtg gta ctt ctg gtt cct 158
Leu Cys Ala Gly Pro Leu His Thr Glu Ala Val Val Leu Leu Val Pro
        15                      20                      25

```


tct gat gat ggg cgt gct ttt ctg ctg cgg agc cgg ctt ctt cat ccg 206
 Ser Asp Asp Gly Arg Ala Phe Leu Leu Arg Ser Arg Leu Leu His Pro
 30 35 40

gag gcg cat gta ccc ccc gcc gct gat cga gga gcc agc ctt caa tgt 254
 Glu Ala His Val Pro Pro Ala Ala Asp Arg Gly Ala Ser Leu Gln Cys
 45 50 55 60

gtc cta cac cag gca gcc ccc aaa tcc cgg ccc agg agc cca gca gcc 302
 Val Leu His Gln Ala Ala Pro Lys Ser Arg Pro Arg Ser Pro Ala Ala
 65 70 75

ggg gcc gcc cta tta cac cga ccc agg agg acc ggg gat gaa ccc tgt 350
 Gly Ala Ala Leu Leu His Arg Pro Arg Arg Thr Gly Asp Glu Pro Cys
 80 85 90

cgg gaa ttc cat ggc aat ggc ttt cca ggt ccc acc caa ctc acc cca 398
 Arg Glu Phe His Gly Asn Gly Phe Pro Gly Pro Thr Gln Leu Thr Pro
 95 100 105

ggg gag tgt ggc ctg ccc gcc ccc tcc agc cta ctg caa cac gcc tcc 446
 Gly Glu Cys Gly Leu Pro Ala Pro Ser Ser Leu Leu Gln His Ala Ser
 110 115 120

gcc ccc gta cga aca ggt agt gaa ggc caa gta gtg ggg tgc cca cgt 494
 Ala Pro Val Arg Thr Gly Ser Glu Gly Gln Val Val Gly Cys Pro Arg
 125 130 135 140

gca aga gga gag aca gga gag ggc ctt tcc ctg gcc ttt ctg tct tcg 542
 Ala Arg Gly Glu Thr Gly Glu Gly Leu Ser Leu Ala Phe Leu Ser Ser
 145 150 155

ttg atg ttc act tcc agg aac ggt ctc gtg ggc tgc taa gggcagttcc 591
 Leu Met Phe Thr Ser Arg Asn Gly Leu Val Gly Cys
 160 165

tctgatatcc tcacagcaag cacagctctc tttcaggctt tccatggagt acaatatatg 651

aactcacact ttgtctcctc tggtgcttct gtttctgacg cagtctgtgc tctcacatgg 711

tagtgtggtg acagtccccg agggctgacg tccttacggt ggcgtgacca gatctacagg 771

agagagactg agaggaagaa ggcagtgctg gaggtgcagg tggcatgtag aggggccagg 831

ccgagcatcc caggcaagca tccttctgcc cgggtattaa taggaagccc catgccgggc 891

ggctcagccg atgaagcagc agccgactga gctgagccca gcaggtcac tcgtccagcc 951

tgctctctcg tcagccttcc tcttccagaa gctgttggag agacattcag gagagagcaa 1011
gccccttgtc atgtttctgt ctctgttcat atcctaaaga tagacttctc ctgcaccgcc 1071
aggggaagggg agcacgtgca gctctcaccg caggatgggg cctagaatca ggcttgcctt 1131
ggaggcctga cagtgatctg acatccacta agcaaattta tttaaattca tgggaaatca 1191
cttctgccc caaactgaga cattgcattt tgtgagctct tggctctgatt tggagaaaagg 1251
actgttacct atttttttgg tgtgtttatg gaagtgcatt tagagcgtcc tgccttttga 1311
aatcagactg ggtgtgtgtc ttccctggac atcactgcct ctccaggga tctcaggcc 1371
cgggggtctc ctccctcag gcagctccag tgggtgggttc tgaaggggtgc tttcaaaacg 1431
gggcacatct ggctgggaag tcacatggac tcttccaggg agagagacca gctgaggcgt 1491
ctctctctga ggttgtgttg ggtctaagcg ggtgtgtgct gggctccaag gaggaggagc 1551
ttgctgggaa aagacaggag aagtactgac tcaactgcac tgaccatgtt gtcataatta 1611
gaataaagaa gaagtggctg gaaatgcaca ttctggata ggaatcacag ctacccccag 1671
gatctcacag gtagtctcct gagtagttga cggctagcgg ggagctagtt ccgccgcata 1731
gttatagtgt tgatgtgtga acgctgacct gtcctgtgtg ctaagagcta tgcagcttag 1791
ctgaggcgcc tagattacta gatgtgctgt atcacgggga atgaggtggg ggtgcttatt 1851
ttttaatgaa ctaatcagag cctcttgaga aattgttact cattgaactg gagcatcaag 1911
acatctcatg gaagtggata cggagtgatt tgggtgtccat gcttttctact ctgaggacat 1971
ttaatcggag aacctcctgg ggaattttgt gggagacact tgggaacaaa acagacaccc 2031
tgggaatgca gttgcaagca cagatgctgc caccagtgtc tctgaccacc ctggtgtgac 2091
tgctgactgc cagcgtggta cctcccatgc tgcaggcctc catctaaatg agacaacaaa 2151
gcacaatgtt cactgtttac aaccaagaca actgogtggg tccaaacact cctcttctc 2211
caggtcattt gttttgcatt tttaatgtct ttatTTTTTg taatgaaaaa gcacactaag 2271
ctgcccctgg aatcgggtgc agctgaatag gcacccaaaa gtccgtgact aaatttcgtt 2331
tgtctTTTTg atagcaaatt atgttaagag acagtgatgg ctagggtca acaattttgt 2391

attcccatgt ttgtgtgaga cagagtttgt tttcccttga acttggttag aattgtgcta 2451
 ctgtgaacgc tgatcctgca tatggaagtc ccgcttcggt gacatttcct ggccattctt 2511
 gtttccattg tgtggatggt gggttgtgcc cacttcctgg agtgagacag ctcttggtgt 2571
 gtagaattcc cggagcgtcc gtggttcaga gtaaacttga agcagatctg tgcattgctt 2631
 tcctctgcaa caattggctc gtttctcttt tttgttctct tttgatagga tcctgtttcc 2691
 tatgtgtgca aaataaaaaat aaatttgggc 2721

<210> 18

<211> 168

<212> PRT

<213> Homo sapiens

<223> clone 11A12 - hypothetical protein DKFZp564K0822

<400> 18

Met	Pro	Leu	Leu	Arg	Gly	Leu	Leu	Trp	Leu	Gln	Val	Leu	Cys	Ala	Gly	1	5	10	15
Pro	Leu	His	Thr	Glu	Ala	Val	Val	Leu	Leu	Val	Pro	Ser	Asp	Asp	Gly	20	25	30	
Arg	Ala	Phe	Leu	Leu	Arg	Ser	Arg	Leu	Leu	His	Pro	Glu	Ala	His	Val	35	40	45	
Pro	Pro	Ala	Ala	Asp	Arg	Gly	Ala	Ser	Leu	Gln	Cys	Val	Leu	His	Gln	50	55	60	
Ala	Ala	Pro	Lys	Ser	Arg	Pro	Arg	Ser	Pro	Ala	Ala	Gly	Ala	Ala	Leu	65	70	75	80
Leu	His	Arg	Pro	Arg	Arg	Thr	Gly	Asp	Glu	Pro	Cys	Arg	Glu	Phe	His	85	90	95	
Gly	Asn	Gly	Phe	Pro	Gly	Pro	Thr	Gln	Leu	Thr	Pro	Gly	Glu	Cys	Gly	100	105	110	
Leu	Pro	Ala	Pro	Ser	Ser	Leu	Leu	Gln	His	Ala	Ser	Ala	Pro	Val	Arg	115	120	125	
Thr	Gly	Ser	Glu	Gly	Gln	Val	Val	Gly	Cys	Pro	Arg	Ala	Arg	Gly	Glu	130	135	140	
Thr	Gly	Glu	Gly	Leu	Ser	Leu	Ala	Phe	Leu	Ser	Ser	Leu	Met	Phe	Thr	145	150	155	160
Ser	Arg	Asn	Gly	Leu	Val	Gly	Cys									165			

<210> 19

<211> 4384

<212> DNA

<213> Homo sapiens

<220>

<223> Clone 14H12 - cDNA FLJ10838 fis, clone
 NT2RP4001274, weakly similar to Human transporter
 protein (gl7) mRNA

<220>

<221> CDS

<222> (534)..(1754)

<400> 19

```

agccgcgcga cgccgcgcgc ttagaacgcc tttccagtac tgctagcagc agcccgacca 60
cgcgttaccg cacgctcgcg cctttccctt gacacggcgg acgccggagg attggggcgg 120
caatttgtct tttccttttt tattaataatt atttttcctg cctgttggtg gatttgggga 180
aattttttgt ttgtttttta tgatttgtat ttgactgaga gaaaccact gaagacgtct 240
gcgtgagaat agagaccacc gaggccgact cgcgggccgc tgcaccacc gcccaaggaca 300
aaaggagccc agcgctacta gctgcacccg attcctccca gtgcttagca tgaagaaggc 360
cgaaatggga cgattcagta tttccccgga tgaagacagc agcagctaca gttccaacag 420
cgacttcaac tactcctacc ccaccaagca agctgctctg aaaagccatt atgcagatgt 480
agatcctgaa aaccagaact ttttacttga atcgaatttg gggaagaaga agt atg      536
                                         Met
                                         1

aaa cag aat tta att ctc ttg aca ttt gtg tca ata ttt tcc ctg tat      584
Lys Gln Asn Leu Ile Leu Leu Thr Phe Val Ser Ile Phe Ser Leu Tyr
          5                      10                      15

tct gtt cat ctc ctt ttg aag act gcc aat gaa gga ggg tct tta tta      632
Ser Val His Leu Leu Leu Lys Thr Ala Asn Glu Gly Gly Ser Leu Leu
          20                      25                      30

tat gaa caa ttg gga tat aag gca ttt gga tta gtt gga aag ctt gca      680
Tyr Glu Gln Leu Gly Tyr Lys Ala Phe Gly Leu Val Gly Lys Leu Ala
          35                      40                      45

gca tct gga tca att aca atg cag aat att gga gct atg tca agc tac      728
Ala Ser Gly Ser Ile Thr Met Gln Asn Ile Gly Ala Met Ser Ser Tyr
          50                      55                      60                      65

```

ctc ttc ata gtg aaa tat gag ttg cct ttg gtg atc cag gca tta acg	776
Leu Phe Ile Val Lys Tyr Glu Leu Pro Leu Val Ile Gln Ala Leu Thr	
70 75 80	
aac att gaa gat aaa act gga ttg tgg tat ctg aac ggg aac tat ttg	824
Asn Ile Glu Asp Lys Thr Gly Leu Trp Tyr Leu Asn Gly Asn Tyr Leu	
85 90 95	
gtt ctg ttg gtg tca ttg gtg gtc att ctt cct ttg tcg ctg ttt aga	872
Val Leu Leu Val Ser Leu Val Val Ile Leu Pro Leu Ser Leu Phe Arg	
100 105 110	
aat tta gga tat ttg gga tat acc agt ggc ctt tcc ttg ttg tgt atg	920
Asn Leu Gly Tyr Leu Gly Tyr Thr Ser Gly Leu Ser Leu Leu Cys Met	
115 120 125	
gtg ttc ttt ctg att gtg gtc att tgc aag aaa ttt cag gtt ccg tgt	968
Val Phe Phe Leu Ile Val Val Ile Cys Lys Lys Phe Gln Val Pro Cys	
130 135 140 145	
cct gtg gaa gct gct ttg ata att aac gaa aca ata aac acc acc tta	1016
Pro Val Glu Ala Ala Leu Ile Ile Asn Glu Thr Ile Asn Thr Thr Leu	
150 155 160	
aca cag cca aca gct ctt gta cct gct ttg tca cgt aac gtg act gaa	1064
Thr Gln Pro Thr Ala Leu Val Pro Ala Leu Ser Arg Asn Val Thr Glu	
165 170 175	
aat gac tct tgc aga cct cac tat ttt att ttc aac tca cag act gtc	1112
Asn Asp Ser Cys Arg Pro His Tyr Phe Ile Phe Asn Ser Gln Thr Val	
180 185 190	
tat gct gtg cca att ctg atc ttt tca ttt gtc tgt cat cct gct gtt	1160
Tyr Ala Val Pro Ile Leu Ile Phe Ser Phe Val Cys His Pro Ala Val	
195 200 205	
ctt ccc atc tat gaa gaa ctg aaa gac cgc agc cgt aga aga atg atg	1208
Leu Pro Ile Tyr Glu Glu Leu Lys Asp Arg Ser Arg Arg Arg Met Met	
210 215 220 225	
aat gtg tcc aag att tca ttt ttt gct atg ttt ctc atg tat ctg ctt	1256
Asn Val Ser Lys Ile Ser Phe Phe Ala Met Phe Leu Met Tyr Leu Leu	
230 235 240	
gcc gcc ctc ttt gga tac cta aca ttt tac gaa cat gtt gag tca gaa	1304
Ala Ala Leu Phe Gly Tyr Leu Thr Phe Tyr Glu His Val Glu Ser Glu	
245 250 255	

ttg ctt cat acc tac tct tct atc ttg gga act gat att ctt ctt ctc 1352
 Leu Leu His Thr Tyr Ser Ser Ile Leu Gly Thr Asp Ile Leu Leu Leu
 260 265 270

att gtc cgt ctg gct gtg tta atg gct gtg acc ctg aca gta cca gta 1400
 Ile Val Arg Leu Ala Val Leu Met Ala Val Thr Leu Thr Val Pro Val
 275 280 285

gtt att ttc cca atc cgg agt tct gta act cac ttg ttg tgt gca tca 1448
 Val Ile Phe Pro Ile Arg Ser Ser Val Thr His Leu Leu Cys Ala Ser
 290 295 300 305

aaa gat ttc agt tgg tgg cgt cat agt ctc att aca gtg tct atc ttg 1496
 Lys Asp Phe Ser Trp Trp Arg His Ser Leu Ile Thr Val Ser Ile Leu
 310 315 320

gca ttt acc aat tta ctt gtc atc ttt gtc cca act att agg gat atc 1544
 Ala Phe Thr Asn Leu Leu Val Ile Phe Val Pro Thr Ile Arg Asp Ile
 325 330 335

ttt ggt ttt att ggt gca tct gca gct tct atg ttg att ttt att ctt 1592
 Phe Gly Phe Ile Gly Ala Ser Ala Ala Ser Met Leu Ile Phe Ile Leu
 340 345 350

cct tct gcc ttc tat atc aag ttg gtg aag aaa gaa cct atg aaa tct 1640
 Pro Ser Ala Phe Tyr Ile Lys Leu Val Lys Lys Glu Pro Met Lys Ser
 355 360 365

gta caa aag att ggg gct ttg ttc ttc ctg tta agt ggt gta ctg gtg 1688
 Val Gln Lys Ile Gly Ala Leu Phe Phe Leu Leu Ser Gly Val Leu Val
 370 375 380 385

atg acc gga agc atg gcc ttg att gtt ttg gat tgg gta cac aat gca 1736
 Met Thr Gly Ser Met Ala Leu Ile Val Leu Asp Trp Val His Asn Ala
 390 395 400

cct gga ggt ggc cat taa ttggcaccac tcaaaactcaa actcagtcca 1784
 Pro Gly Gly Gly His
 405

tctgatgccca gtgttgagta aactcaacta ctatgaaatt tcacctaattg ttttcagttt 1844

cacttccttt tgaagtgcag attcctcgct gggtcttctg agtgcagaat aagtgaactt 1904

ttttgttttg ttttgttttt ttaagaaact tatctgtatg ttagaaatgg atatgaacaa 1964

caaaaccacg agtctcgggt taagggaagt gacaatttta ttccattcca gagaatggac 2024

aaactcttaa cttttatcaa gccacatgct tggctgtgtc attgtttaac ttggatatatt 2084
tatgatttta cttgaatgtg cctaattggaa ccatttgatg tgagaaacaa ttcttttttaa 2144
tttacagcaa aatattgaat aaccattgac aaaaacacta ttatttttttg taccaaaaat 2204
acttaaagac ctcagaagca ctcttttact tttaagaaat tgcttttttg aactttattc 2264
agaagcagtt atcaataaat tccataaaat aatgtcattg gtatttataa atgaatatta 2324
atataatgaa atggtttgcc tttttgtagg cataataagc caaatacttt ttaccocaaa 2384
ataattttta gagaaaatga tgtaatgaaa aattgtacca tgaattagga gcatagtttt 2444
ttccatttaa acgtcaccat tacttaaaag atgattgatt gttgctatac caaatcagat 2504
gaactctggt catcactttt cttctctgtc cccaaacaat ttggttcatt cagactgaaa 2564
tgtttgtgtc ttcaacttat tagaatggaa gataatgcag atatttctgt gggaaataaa 2624
ataactaatt ttgagggtacc aaatagtgc aattgggtaaa acagggttta ttcagttgca 2684
tctgtctcca gtgttggtatt gacagctctg ggtctttttt ttgggccagc ccttttttga 2744
cattgcttcc agcagtggaa aatgggcatt tgatggcaat aggccaaaat tattgtgtcc 2804
agggagtaca ctttttcaaa atgctcacct actggaagtg tgaattactt gacaatgtat 2864
ggcttagttg tgttcatgtt ttgtctacag tagagggtcta atccacaggt tacacctatg 2924
ttgatatga tataagttct ctttgcgtag gccactgggt ttctcatgca gtaagcttta 2984
taaaaactca tttgcactgg actgtcatct cattcttgta caacgtagaa ttacttgttt 3044
acatccaaca aatgggttagc tagggaaaac agtgcaaaact gagtgtagt agtcattttg 3104
gtccaactgc atgtcaaccc ttccatttca atccagttta gaaatgaaaa taattacttt 3164
gaaacttggc tttaagagca catttatcgt acgtcacagt gtatggtgaa tatattatta 3224
aataatgtgg tacttcgctc atcaggcata atgtctaaaa tctaataatac ataattccat 3284
taagtgggtg aaggaagcaa ataatggaat tgtcaattgg tcatctggct gtaaggtttg 3344
cccttgaact aaaaatgttg tttggggcaa gggccagaaa tgtggagaca tggtttttgt 3404
tacgcattct tgtattatat gtgactaaat ttacaaacaa gatacatgtg taattaaaga 3464

```

cccttatgga actggaagac gtcttgtagt gctacattgg gtgaaaccgt tgggccattt 3524
ttgtctgttt ctatgaagat aaaataattg ggggccatct agaaatagaa aggcagtggg 3584
aagacagatt ctacggcact gctttcattt aattgggctt taggcactcc attcgaatgc 3644
agaacctcac ctctagttga gaccaagaat tggcaaattt gcatgagctc ctggaaagag 3704
ttgctgactt tgtatctaag acctgccagg gaataccaag agttgtttct acagactttt 3764
tttttttttt gtatgggaga agatactgtg gcaaccagga aggaatggaa aaaaaattct 3824
tttctctaca gcaaattaat gtgaggaagc tcctccaatc ctctggctat ttaaggttca 3884
aatcaagtg cctagggaaa attccaatgg atgattttct gggagctatc ttgtctacct 3944
tgaggttcct gaacaatgaa ttcccattaa tgagcagtct tcagtattaa aaccactgtc 4004
ttgtcacctc attttgcatt actgtcttcc gtggatgttt cagttacaac tgtaatgtta 4064
tttatagaac aacattaatc cattaaagct aacctatttt tcaatattta tgataatcta 4124
tgtacatata ttgtctgtcc atatgtattt gtaaataagg tgtatataat gtcaggtttg 4184
ggctcttgggt tcaagtgtat atattcctgt aagtttctta actgcatttt gatgaattca 4244
cattatgtaa ctataagaat tgtcccaaaa gtacctgtac agaaaattga atattgaaaa 4304
attgacaaat tgtgtacaaa cactaaaaaa aacttgttta aattgtattt gcaataaaca 4364
acatcaaatt ttttcatgac 4384

```

<210> 20

<211> 406

<212> PRT

<213> Homo sapiens

<223> Clone 14H12 - cDNA FLJ10838 fis, clone
NT2RP4001274, weakly similar to Human transporter
protein (g17) mRNA

<400> 20

```

Met Lys Gln Asn Leu Ile Leu Leu Thr Phe Val Ser Ile Phe Ser Leu
  1             5             10             15
Tyr Ser Val His Leu Leu Leu Lys Thr Ala Asn Glu Gly Gly Ser Leu
          20             25             30
Leu Tyr Glu Gln Leu Gly Tyr Lys Ala Phe Gly Leu Val Gly Lys Leu
        35             40             45

```


Ala Ala Ser Gly Ser Ile Thr Met Gln Asn Ile Gly Ala Met Ser Ser
 50 55 60
 Tyr Leu Phe Ile Val Lys Tyr Glu Leu Pro Leu Val Ile Gln Ala Leu
 65 70 75 80
 Thr Asn Ile Glu Asp Lys Thr Gly Leu Trp Tyr Leu Asn Gly Asn Tyr
 85 90 95
 Leu Val Leu Leu Val Ser Leu Val Val Ile Leu Pro Leu Ser Leu Phe
 100 105 110
 Arg Asn Leu Gly Tyr Leu Gly Tyr Thr Ser Gly Leu Ser Leu Leu Cys
 115 120 125
 Met Val Phe Phe Leu Ile Val Val Ile Cys Lys Lys Phe Gln Val Pro
 130 135 140
 Cys Pro Val Glu Ala Ala Leu Ile Ile Asn Glu Thr Ile Asn Thr Thr
 145 150 155 160
 Leu Thr Gln Pro Thr Ala Leu Val Pro Ala Leu Ser Arg Asn Val Thr
 165 170 175
 Glu Asn Asp Ser Cys Arg Pro His Tyr Phe Ile Phe Asn Ser Gln Thr
 180 185 190
 Val Tyr Ala Val Pro Ile Leu Ile Phe Ser Phe Val Cys His Pro Ala
 195 200 205
 Val Leu Pro Ile Tyr Glu Glu Leu Lys Asp Arg Ser Arg Arg Arg Met
 210 215 220
 Met Asn Val Ser Lys Ile Ser Phe Phe Ala Met Phe Leu Met Tyr Leu
 225 230 235 240
 Leu Ala Ala Leu Phe Gly Tyr Leu Thr Phe Tyr Glu His Val Glu Ser
 245 250 255
 Glu Leu Leu His Thr Tyr Ser Ser Ile Leu Gly Thr Asp Ile Leu Leu
 260 265 270
 Leu Ile Val Arg Leu Ala Val Leu Met Ala Val Thr Leu Thr Val Pro
 275 280 285
 Val Val Ile Phe Pro Ile Arg Ser Ser Val Thr His Leu Leu Cys Ala
 290 295 300
 Ser Lys Asp Phe Ser Trp Trp Arg His Ser Leu Ile Thr Val Ser Ile
 305 310 315 320
 Leu Ala Phe Thr Asn Leu Leu Val Ile Phe Val Pro Thr Ile Arg Asp
 325 330 335
 Ile Phe Gly Phe Ile Gly Ala Ser Ala Ala Ser Met Leu Ile Phe Ile
 340 345 350
 Leu Pro Ser Ala Phe Tyr Ile Lys Leu Val Lys Lys Glu Pro Met Lys
 355 360 365
 Ser Val Gln Lys Ile Gly Ala Leu Phe Phe Leu Leu Ser Gly Val Leu
 370 375 380
 Val Met Thr Gly Ser Met Ala Leu Ile Val Leu Asp Trp Val His Asn
 385 390 395 400
 Ala Pro Gly Gly Gly His
 405

<210> 21
 <211> 588
 <212> DNA
 <213> Homo sapiens

<220>

<223> clone 1F2 - hypothetical protein HSPC014 (HSPC014)

<400> 21
 tctcttgnaa gctccccaaa aaggccaacg ttgcgagcgc tgcggaagat gaatgccaga 60
 ggacttggat ctgagctaan ggacagtatt ccagttactg aactttcanc aagtggncct 120
 tttgaaagtc atgatcttct tnnngaaangn ttntctngtg nnaaaaaatga acttttgcct 180
 agtcatcccc ttgaattatc agaaaaaaaaat ttccagctca accaanataa aatgaatttt 240
 tccacactga gaaacattca gggtnatatt gctccgctaa aattacagat ggaattcaag 300
 gnagtgcagc angttcagcg tcttccattt ctttcaagct caaatctttc actggatgtt 360
 ttgaggggta atgatgagac tattggattt gaggatatnn tcannngntnc ctctcnaagc 420
 gaagncatgg gagagacaca cttgntggng gaatntnanc tcggctnact gtgatagnga 480
 gctnggtcat ggaaaccgag ggntgaacnt ggnaatagnn atcttggccc tgnantcgct 540
 gcaccacanc cttnaaagtc ctggcncccc tnaaaaaaaaa naaaaaaaa 588

<210> 22
 <211> 815
 <212> DNA
 <213> Homo sapiens

<220>

<223> clone 2H4 - hypothetical protein MGC2577
 (MGC2577)

<400> 22
 cccgaccccc caagcccact ggtgaaacag ctgagtgaag tatttgantt ctgaagactc 60
 taaatcaaat cttccccag ngctgtttt gccccagag gcacctttat cttctgaatt 120
 ggacttgctt ctgggtaccc agttatctgt tgaggaacag atgccacctt ggaaccagac 180
 tgagttcccc tccaaacagg tgttttccaa ggaggaagca agacagccca cagaaacccc 240
 tgtggccagc cagagctccg acaagccctc aaggggaccct gagactccca gatcttcagg 300
 ttctatgcgc aatagatgga aaccaaacag cagcaaggta ctaggagat cccccctcac 360
 catcctgcag gatgacaact cccctggcac cctgacacta cgacagggtta agcggccttc 420
 acccctaagt gaaaatgtta gtgaactaaa ggaaggagcc attcttggaa ctggacgact 480
 tctgaaaact ggaggacgag catgggagca aggccaggac catgacaagg aaaatcagca 540
 ctttcccttg gtggagagct aggccttgca tggccccagc aatgcagtca ccagggcct 600
 ggtgatattt gtgtcctctc accccttctt tcccagggat actgaggaat ggcttgtttt 660
 cttagacttc tntcagcta ccaaactgta ctgcacccat tttggtggac gatgaaatgg 720
 aaatagcccc taatatgtca aagccaaaaa tacccttttt gaaaagccgc cttggaagtc 780
 nngaataact tcnttcacaa ctggatgatg gttac 815

<210> 23
 <211> 533
 <212> DNA
 <213> Homo sapiens

<220>
 <223> clone 3C2 - clone FLC0593

<400> 23
 ggccaaccgg cgcgtttttt tttctgctta aaaatttcaa ttctcgtggt aataccagag 60
 tagaaggaga gggtagacttt accgaactgn nggccattgg ggaggcagat gcgggtntgg 120
 aggtgtgggc tgaaggnant gactgtttga ttttaaaaag tgtgactgtc agttgtatct 180
 gttgcttttc tcaatgattc agggatacaa atgggcttct ctcatcatt aaaagaaaac 240
 gcgacatctt tctaagattc tctgtgggaa aatgactgtc aataaaatgc nggtttcttg 300
 gccattcgtc ttactttcat tttttgatta caaatttctc ttgacgcaca caattatgtc 360
 tgctaatcct cttcttccta gagagagaaa ctgtgtcctt tcagtgttgc tgccataaag 420
 gggtnngggg aatccattgt aaaagtccca ngttctaaat taactaaatg tgtacanaaa 480
 tgaacgtgta agtaatggtt ctacagggtt ttgcaacaaa actggccttt cgt 533

<210> 24
 <211> 547
 <212> DNA
 <213> Homo sapiens

<220>
 <223> clone 3F11 - BAC clone CTD-2314H8

<400> 24
 gaagaactng ttctacaatt cagctgtatg ttgtaaagga gtcaagtcaa tgaaggagtt 60
 ataaagagtt gctgaaaaat ggagnaacag tgtatgggac ttcacatcac ntctccttat 120
 tgattgttan nttgatccct cttgttcttt tgttgtaaac attttctata attaggaaat 180
 gccatttaag agtgagagag gtatatatct atgagccatt gtgtttggtg ttttacaaga 240
 actttacat actggtgtgt agtccattct gtacagttta aaagtgattc acgatttgca 300
 ggctttttat cagatcacia aaaaatcagt ctttaagcat ttgcttggtg aggtttctta 360
 agattaggtt tataatataa ccatctgtaa tgtatctctc gtttgagctt gtgggccata 420
 caattcatta actagatgaa tacattgtgg acagcatcct cactaccctt ctctactcac 480
 tcacaaagaa ccatgataca ctggaatgtt tttctctgga atcctctttc tactctttna 540
 ttaaaat 547

<210> 25
 <211> 537
 <212> DNA
 <213> Homo sapiens

<220>
 <223> clone 4D10 - full length insert cDNA clone ZE03C06

<400> 25

```

gcttggccna aaaggccaac taaaagccta cgtgttggt aatccaacag caagggagat 60
ttttgaatca taataactca taaggtgcta tctgggnagt gatgccctca gaggnnttgc 120
tgttagctgg cagctgacgc tgctaggata gttagtttgg aaatgggtact tcataataaaa 180
ctacacaagg aaagtcagcc accgtgtctt atgaggaatt ggacctata aatttttagtg 240
tgccttccaa acctgagaat atatgctttt ggaagttaaa atttaaattgg cttttgccac 300
atacatagat cttcatgatg tgtgagtgtg attccatgtg gatatcagtt accaaacatt 360
acaaaaaaaa tttatggccc aaaatgacca acgaaattgt tacaatagaa tttatccaat 420
tttgatcttt ttatatctt ctaccacacc tggaaacaga ccaatagaca ttttgggggtt 480
ttataatggg aatttgtata aagcattact ctttttcaat aaattgtttt ttaattt 537

```

<210> 26

<211> 582

<212> DNA

<213> Homo sapiens

<220>

<223> clone 4E9 - hypothetical protein dJ465N24.2.1
(DJ465N24.2.1)

<400> 26

```

tggccaaaaa ggccaaccct gtctagctgt cggaaaagg aacagaagat ggaactcgaa 60
atcccaatga ngnacctacc cagcaaagan gggtagcttt tagctctaata aggttctgta 120
gcaaagccaa taaaaaaatc agctaaagct gccacagaag aggcattctc aagatcacca 180
aaaatagatc agaaaaaaag tccatatgga ctgtggatac ctatctaaaa gaagaaaact 240
gatggctaag tttgcatgaa aactgcactt tattgcaagt tagtgtttct agcattatcc 300
catccctttg agccattcag gggtaacttg gcatttaaaa accaacacaa aaagatgtaa 360
atacttaaca ctcaaattt aacatttttag gtttctcttg cagatatgag agatagcaca 420
gatggaccaa aggttatgca cagggtgggag tcttttgtat atagttgtaa atattgtctt 480
ggttatgtaa aaatgaaatt ttttagacac agtaattgaa ctgtattcct gttttgtata 540
tttaataaat ttcttgtttt cattcttaaa aaaaaaaaaa aa 582

```

<210> 27

<211> 388

<212> DNA

<213> Homo sapiens

<220>

<223> clone 4F7 - chromosome 1 clone RP11-109I2

<400> 27

```

tttttttncc ctggatcagc tcggcctnna aggcctcgga aaacaagatg cttcctctgg 60
aatgtgagtc caaagagtta ccagcgctgc cctctagtga tctcagctca ggntatgcac 120
taaccgtgtg gntacagggc tgagtagtgc tgcagtgtga agtgaatgga aggcctcgag 180
gtgtttgtgg ctggccaccc tgatcagcct gcaggtagtc ccgatgaagc cagggcacag 240

```

```

ggggattcgt tccagcttgt tcactttatt ctgccttgcc aggttactga aagtccctcg 300
tttgctctca ccagccttcc tggaaatgtg gactcttgaa agaaaagctc cegtgcctctt 360
gaagtatacc tgcttgccan gggagtc 388

```

<210> 28

<211> 605

<212> DNA

<213> Homo sapiens

<220>

<223> clone 4S-10d4 -DNA sequence from clone RP1-64K7 on chromosome 20q11.21-11.23 Contains the EIF2S2 gene for eukaryotic translation initiation factor 2 subunit 2 (beta, 38kD), a putative novel gene, the

<220>

<223> ribonucleoprotein RALY or autoantigen P542, an RPS2 (RPS4) (40S ribosomal protein S2) pseudogene, ESTs, STS, GSSs and two CpG islands

<400> 28

```

tgcttggtca tgctggcacc gggcatatg ctggacaggg agaacgagag tcccatcctg. 60
gaaatccaga aaagccctg gatgctccag cccctgggaa agcacacagc caggcccttg 120
ggtgggaggt tggcttctaa cagtgcatac acatgccctt cctctgagtc ggggcagcaa 180
aaacatccat tccgctgcgc aacagttgtc atttttctaa catctgaaaa ctccagaagg 240
agatggtgat aaatgtggta ccggattctg cctaaaggat cagtcttttag atgttttcag 300
attgaaagcc tcatttgtga tcctcacagc catcttgaaa gaatagagca gccagtggg 360
atactggatt gtgagctaag aggctggga ctttccccct gttgctgcca gccaggttga 420
tgaccctggg caagtctttt tccttaccag gtctcagttt cctcagctgt aaaatgagag 480
gttgatctgg atcagggata gtaaattgggc ctttgttcag ttactgactg ttgtataaca 540
aaccaccccc aaatttagta gccttaataa acatttatta gctcctgaaa aaaaaaaaaa 600
aaaaa 605

```

<210> 29

<211> 661

<212> DNA

<213> Homo sapiens

<220>

<223> clone 4S-10G6 - similar to C50F4.16.p (LOC256281)

<400> 29

```

tgggggtgagt ggccaagact ggcctctgtc tagaaccctg gagtctcact ggagatccag 60
gttggggggcc acctggctga ggaaccatga gacaccaaag atgacgccga gggctctggg 120
gatgtccggg tcgtctgcaa aggcctgggc gccttcagg ggcaggtgca ccacctcaat 180

```

```

gagctcacc ctcctccacca ggccccccacc tggaccgcta cgctgggcat ctgtcacctc 240
tgtgtagaac atgggtctgtc tggagccagt cagtcccact ccagaccagt atgtggcgac 300
ccggcgcgaga tcagagggggg ccaagtggta gccacactcc tcccaagcct ccttgcaagc 360
cacttcctcc agcgagagacc caggctgggtc cacgaggccg gcacacagct caactgtcac 420
ccccgctgag ccgggcagggt ctggctgttag ctcccgaggc ccgtcctgggt ctacagctgc 480
tagggaccct gggaagcggc gctccacctc acccgcatat acagctggcc ggaactgctt 540
caccaacacc aggtctcctcc gagaagagtt gaataagaga acggtcacgc tgtcatgcgt 600
cttcatgaag tcccaggact tctgggcacc attctggcgg taatgcagcg tgagcggccg 660
c

```

<210> 30

<211> 667

<212> DNA

<213> Homo sapiens

<220>

<223> clone 4S-11B8 - DNA sequence from clone RP11-735A5
on chromosome 1

<400> 30

```

tttttttttt tttttttaaa gaaaacctac tgtatgccat ttgcaagaga tataacaaaa 60
gcaaaactgat atgaaaaggc tgaaataaag ggatagttgg tttctggaaa aactaacaac 120
aacaacaaaa atcttccatg aatttttgtt gttgttattt taattttttt tgagacacgg 180
tcttgctgtc acccaggctg gagtacaggg gcacaatcat ggctcacggc agcatcgacc 240
tcctaggctc aaacgatcct ccacttttag cttcccaa at agctggaact acaggtgtgc 300
accgccacgc ctggctactt aaattttttt tttttttaat ttgtacagac atctcaccat 360
gttgcccaag ctggtctcca gtccttgggc tcaagtgatt caccacactg gcactctcaa 420
agtgtgggga ttataggcat gagccactgt actcggcatt ccactgatac ttgataatga 480
aaaaaaagag tottaatacc caatattaga aatacaaaga gaaacagaag aacaaatcag 540
caaaggttta aaagaacaaa aattttatat atatatat ataaacaaat tgatctcaat 600
aaataacaaa ataaacaaaa tggacaaact cttaaaaaa tataccttac caaaactacc 660
tcaaaaaa

```

<210> 31

<211> 578

<212> DNA

<213> Homo sapiens

<220>

<223> clone 4S-11H3 - cDNA FLJ14279 fis

<400> 31

```

ttgtcaaaaa tccattcctg tcccctccac cctgttttta ttctctatc cactggaaac 60
cattaaaaaa aaaagttttt gatttatagt caattcttgt atttgaggta gttgtgttct 120
ataaagtgtg gcaaatactg aattagcaaa taccaaatta ttactactag ggcaaataca 180
cgattaagtt tctgtgagcc tctggtaata tttttgtcaa ccaatcgata catttacttt 240

```

```

gttttatgag tgtttctgtt aaagatacct tatttagtat atattgttga ttcattaacg 300
tttaactcac tgccaatagt attaaaaactc atatgaatga aatgtgtcta acacgtattt 360
tccttataag tcatatcatg gccttcttga gcttaagaac actagagagc acttcagcac 420
tacacttgag ggccatttta ttactttatt ttacagatg gatcttgcta tattgcccac 480
actggactgt aatggctatt cacagggtgcc aaagtagcat actgcagact caaactcctg 540
gcctcaagcc atccacctca tcctcccaag tagttggg 578

```

<210> 32

<211> 352

<212> DNA

<213> Homo sapiens

<220>

<223> clone 4S-12A9 - chromosome 18, clone RP11-13N13

<400> 32

```

acatcacttg aggtcagtag acagggtttc accatgttac ccaggctggt ctggaactcc 60
tgacctcaag tgattcgccc atctcgccct gccaaagtgc tgaggtcaca ggcatgaggt 120
caggagaacg agaccatcct aacacagcga aaccccgctc ctactaaaaa gacaaaaaat 180
tagctgggtg tgggtggcagg cacctgtagt cccagctact cgggaggctg aggcaggaga 240
atggcctgaa cccgggaggc agagcttgca gtgagccaag atcgcgccac tgtactccag 300
cctgggcaac agagcaagac tgtctccaga aaaaaaaaaa aaaaaaaaaa aa 352

```

<210> 33

<211> 469

<212> DNA

<213> Homo sapiens

<220>

<223> clone 4S-12E6- hypothetical protein MGC955

<400> 33

```

gggagagtgg tcagcagtat ttttccatca tttctccaaa ggaatggggg acaagttgtc 60
cacatgactt ccttgggtgcc tacaaactac agcatgactt gtcctggact ccgtatgagg 120
acattgagaa gcaagatgct aaaatcagca tgatggacat gttgctaagc cagtcagtgg 180
ccctgcctcc gtgcactgaa cccaacttcc agggactgac tcaactgagag tgggctttga 240
caaacagctc tcacaggacc tggctgtcaa cctccttggt gccccactg ttgccttgag 300
aattgaagac atgtaggtga ctacaaaact tcttggaaag agaccctgtg tgaatgtaaa 360
tgctgtcatt atgactttta attgggatgg gaataatcat tgagacagag tcaactgtctt 420
tcgggatcct ctttggacca cagataccca agtcagtcag tttcagagt 469

```

<210> 34

<211> 795

<212> DNA

<213> Homo sapiens

<220>

<223> clone 4S- 14D4 - clone IMAGE; 4778940 mRNA

<400> 34

```

aaaagcagat ttaaaaagca gctcacagtt gggcgcggtg gctcacacct gtaatcccag 60
cactttggga ggccgaggcg ggtggatcac cagagggtcag aagttcgaga ccagcctggc 120
caacgttggtg aaaccctgtc tctactaaaa atacaaaaat tggccacgtg tgggtggcggg 180
cacctgtaat cctagctact tgggaagctg agacaggaga attgcttgaa ccgaggaggt 240
ggagtgcagt ggcatgatta aggttcactg cagcctcaat ctcccactct ccagcgatca 300
tcccacctca gcctcttggg tagctgggac cacaggcacg agctaccatg cctggctaata 360
ttattttttg tagagacggg gtttcgccat gttgcccagg ctggtctgga atttctgacc 420
tctggcagtc cacctgtttc cgcctcccaa atgctgagat tagaggcatg agtcactgca 480
cccagcccg cgcctctttt ataaggcat taatcccctt catgagggat attctctcat 540
gacttaata cctgccaaag accccacctc ttaatactac attaagtgat ggggtcaatg 600
tattnaattg gggggggagg gggcacattc agaccatagc atcttagtca ttcttggttt 660
tatttaagat tatttagact ganggcattg aaaaatagca tacttggatg ggacttcagc 720
attcnatnag tggccttaat aaaacctggt aattaaaaaa gcncttaatt ttgcaaaaaa 780
aaaaaaaaa aaaaaa

```

795

<210> 35

<211> 324

<212> DNA

<213> Homo sapiens

<220>

<223> clone 4S-2A6 - PAC clone RP3-414A15 from 14q24.3

<400> 35

```

tggcgtgaac ccagaggtg gagcttgtag gaggtgacag cgtgctggca gccctcacag 60
cacttctct gcctgggctc ccactttggc ggcacttgag aagcccttca gccaccgct 120
gcactgtggg agcccctttc tgggctggcc aaggccggag ccgtctccct cagcttgtag 180
ggaggtgtgg agggagaggc gcaagtggga actgaggctg cgttccgcac ttgcgggcca 240
gctggagtgc cagggtgggca tgggcttggt gggccccaca ctcgagagccg ccagctggcc 300
ctgccagccc cgggcaatga gggg

```

324

<210> 36

<211> 226

<212> DNA

<213> Homo sapiens

<220>

<223> clone 4S-2D3 - DNA sequence from clone RP11-357H24
on chromosome 10

<400> 36


```

gtctgctgcc tactcgtatg taatatgtgt acataaaagc ggcagctggg ttttcgttta 60
agagtaatct aatatacaga atttgggccc ttaagggttt atacctcttc atttaaaatg 120
ctttctggac aatctgctac caaacacatt ttgttatagg tgacattaaa actacatata 180
aatctacctg cacgacaaca cataaaaaaa aaaaaaaaaa aaaaaa 226

```

<210> 37

<211> 377

<212> DNA

<213> Homo sapiens

<220>

<223> clone 4S-2E2 - cDNA FLJ14385 fis, clone
HEMBA1002212, weakly similar to TYROSINE-PROTEIN
KINASE 2

<400> 37

```

gttggccttt gtttaaaaca ctgaaccttt tgctgatgtg tttatcaaata gataactgga 60
agctgaggag aatatgcctn aaaaagagta gtccttgga tacttcagac tctgggttaca 120
gattgtcttg atctcttgga tctcctcaga tctttgggtt ttgctttaat ttattaaatg 180
tattttccat actgagttta aaatttatta atttgtacct taagcatttc ccagctgtgt 240
aaaaacaata aaactcaaata aggatgataa agaataaagg acactttggg taccagaaaa 300
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 360
aaaaaaaaaa aaaaaaa 377

```

<210> 38

<211> 758

<212> DNA

<213> Homo sapiens

<220>

<223> clone 4S-2F8 - chromosome 17, clone hRPC.1110_E_20

<400> 38

```

caaaaaaaaaa aaaaaaaaaa aaaaaacatc agatcttgtg agacttattc actatcagga 60
gaccagcaca cgaggaaagac cctcatgatt caattacctc ccaccagggt cctcccacaa 120
cacataggaa ttatgggagc tgcaattcaa gatgacaatt ggggtggggac acagcaaaac 180
cacatcacat gctgagctgt agcagggtgaa taaaccactg agactacgaa cctctgtcct 240
ctgaagaaga tgccattttc tcaaattcta ggagtgtggg cactgttctt gaggttgctg 300
cagaggagca ctgctggaga aaaggaaggg ggaaatccac atatccacta agacatcgctc 360
agaatactgc acatggaatt ataggatatt aacaatttaa gagactgtag acacaggata 420
gtcaaaatct cccattttac agacaaatca acagaggccc agagcagtga aggcatctac 480
ccagacactc actggatcgg tgcagctgga tctagatcca ggtctcttga ctcatttaac 540
agctgttaaa ccaaaaatgg gtgtgattta gtcccattgt catctgatac attggcaatg 600
ccctgcataa ttttttgtct ctatgtttta tactcctttg taaagggtag cttttgatata 660
ttcctgaact gagcatctgt taaaattgga ttcactcttc cttaaggca acaattgggtg 720
tttctgatct ttaatgccaa aaaaaaaaaa aaagaaaa 758

```

<210> 39
<211> 840
<212> DNA
<213> Homo sapiens

<220>
<223> clone 4S-3C2 - clone FLC0593

<400> 39
tgactttacc gaactgacag ccattgggga ggcagatgcg ggtgtggagg tgtgggctga 60
aggtagtgac tgtttgattt taaaaagtgt gactgtcagt tgtatctgtt gctttttctca 120
atgattcagg gatacaaatg ggcttctctc attcattaaa agaaaacgcg acatctttct 180
aagattctct gtgggaaaat gactgtcaat aaaatgcggg tttctgggcc attcgtctta 240
ctttcatttt ttgattacaa atttctcttg acgcacacaa ttatgtctgc taatcctctt 300
cttcctagag agagaaaactg tgctccttca gtgttgctgc cataaagggg tttggggaat 360
cgattgtaaa agtcccaggt tctaaattaa ctaaatgtgt acagaaatga acgtgtaagt 420
aatgtttcta caggctcttg caacaaactg tcactttcgt ctccagcaga gggagctgta 480
ggaatagtgc ttccagatgt ggtctcccggt gtggggccca gcaatggggg cccctgatgc 540
caagagctct ggaggttctt gaaagagggg acacgaagga ggagtgactg ggaagcctcc 600
catgccaaagg aggtgggagg tgccctggaa atagctgcct catgccactt aggccatgac 660
tggatttaaat gtcagtgggt tgccacagtg cagtggctag acaactgaaa ggggctacca 720
aggctgggaa aaaaatgcaa ttgttgctgt gagtgacttt gaaagactct ggtgccttgt 780
ggtgcccttc tgaaattcaa acagtaatgc aaaagtgtct gcattagaat ttacgggtgc 840

<210> 40
<211> 480
<212> DNA
<213> Homo sapiens

<220>
<223> clone 4S-3F11 - BAC clone CTD-2314H8

<400> 40
agagttgctg aaaaatggag taacagtgtg tgggaacttca catcagttct ccttattgat 60
tgttagtttg atccctcttg ttcttttgtt gtaaacattt tctataatta ggaaatgcc 120
tttaagagtg agagaggtat atatctatga gccatttgtt ttggtgtttt acaagaactt 180
taccatactg gtgtgtagtc cattctgtac agttttaaag tgattcacga tttgcaggct 240
ttttatcaga tcacaaaaaa atcagtcctt aagcatttgc ttggttaagg ttcttaagat 300
taggtttata atacaacat ctgtaatgta tctctcgttt gagcttgtgg gccatacaat 360
tcattaacta gatgaatata ttgtggacag catcctcact acccctctct actcactcac 420
aaagaacat gatacactgg aatgtttttc tctggaatcc tctttctact cttgtattaa 480

<210> 41
<211> 506

<212> DNA

<213> Homo sapiens

<220>

<223> clone 4S-4C11 - similar to putative, clone
MGC:33177 IMAGE:4823662

<400> 41

```
atcatttgat aatttacctt agagcattta aaaaaatata atcaaactaa ttgccagcca 60
agtcagtcac cctcctggga gtatatagag tccaaggtt agcgctcctg tattagacta 120
tttcaatttt aggaaaatca tgaccatgtg gggaaacaat gactttaaaa tgctgaaatt 180
aaaatttatg ctttaactgg aatatttttt gcttaactac tcaattagaa tattgtacac 240
ctgatcaatg tgtgttcagc acagatggcc atgaattgtc atttatagtc caatttttta 300
tcttaatcat aaaatgttta ggaatctatg aaatttaact ttaggaacaa aacgttttagc 360
agggttgatt gatattatgt ttacattgtt ctggcaatcc acagaaagag aagagcctta 420
atttttaaaa cccatttttag tcattttatg acaattaaag ttgtttaata aacatctttt 480
ttcaaagaag caaaaaaaaa aaaaaa                                     506
```

<210> 42

<211> 558

<212> DNA

<213> Homo sapiens

<220>

<223> clone 4S-4E9 - hypothetical protein dJ465N24.2.1
(DJ465N24.2.1)

<400> 42

```
ctgtcggaaa aggtaacaga agatggaact cgaaatccca atggaaaacc taccagcaa 60
agaagcatag cttttagctc taataattct gtagcaaagc caatacaaaa atcagctaaa 120
gctgccacag aagaggcatc ttcaagatca caaaaaatag atcagaaaaa aagtccatat 180
ggactgtgga tacctatcta aaagaagaaa actgatggct aagtttgcac gaaaactgca 240
ctttattgca agttagtgtt tctagcatta tccatccct ttgagccatt caggggtact 300
tgtgcattta aaaaccaaca caaaaagatg taaatactta acactcaaat attaacattt 360
taggtttctc ttgcagatat gagagatagc acagatggac caaagggttat gcacaggtgg 420
gagtcttttg tatatagttg taaatattgt cttgggttatg taaaaatgaa atttttttaga 480
cacagtaatt gaactgtatt cctgttttgt atatttaata aatttcttgt ttccattctt 540
aaaaaaaaaa aaaaaaaaaa                                     558
```

<210> 43

<211> 352

<212> DNA

<213> Homo sapiens

<220>

<223> clone 4S-4F7 - chromosome 1 clone RP11-109I2

<400> 43

```

gggaaaacaa gatgcttcct ctggaatgtg agtccaaaga gttaccagcg ctgccctcta 60
gtgatctcag ctcagcatat gcactaaccg tgtgtttaca gggctgagta gtgctgcagt 120
gtgaagtga tgaagggcct cgaggtgttt gtggctggcc accctgatca gcctgcaggt 180
agtcccgatg aagccagggc acagggggat tctgtccagc ttgttcactt tattctgcct 240
tgccagggtta ctgaaagtcc ctctgtttgct ctcaccagcc ttcctggaaa tgtggactct 300
tgaaagaaaa gctcccgctg tcttgaagta tacctgcttg ccaggggagt cc 352

```

<210> 44

<211> 524

<212> DNA

<213> Homo sapiens

<220>

<223> clone 4S-5C4 - similar to KIAA0674 protein

<400> 44

```

cccttcctta agcatgattt tgcacagcca accctgggtc taggcgaacc acagggtgag 60
gtcaagggtga gcattctggg aacaatattt gggctcagag ggtgggttgg ccaccttctg 120
agccccaccc cgcagagacc tggatgaagag gatcataacc ctgtcttcaa gaacactggg 180
atttcagcag caagttggaa gaaggactgg taggttcccc tccaagccag tcacctgtaa 240
gagtctgtgc ctctgccaga ctttttaatc tcttcattaa ctctcagact gacctgggag 300
ccctcctcta cctgaatcca gtgctcaact gtgccccggc aacaagacct gggctgaggt 360
ctccctggta gaactaaggg agattacacc atctaaatcc cagtgcagtc aacagcctgg 420
cctatagtcc tgggacatgt atcttcttct ttgccttaaa tctgatacaa gaggtcaatg 480
actttgaaaa taaaactaaa ataaatgtca aaaaaaaaaa aaaa 524

```

<210> 45

<211> 891

<212> DNA

<213> Homo sapiens

<220>

<223> clone 4S-6F6 - similar to RIKEN cDNA 1110012M11 gene

<400> 45

```

atggattcca aggcctctgt gtccaagaag aaacgcatgt gtgtgaagct gttgcccttg 60
ggagccacgg acacggctgt gtttgatgtc cggctgagtg ggaagaccaa gacagtgcct 120
ggataccttc gaatagggga catgggcggc ttgcatctt ggtgcaagaa ggccaaggcc 180
ccgaggccag tgcccaagcc ccgaggtctc agccgggaca tgcagggcct ctctctggat 240
gcagccagcc agccaagtaa gggcggcctc ctggagcggg cagcgtcaag gctgggctct 300
cgggcatcca ctctgcggag gaatgactcc atctacgagg cctccagcct ctatggcatc 360
tcagccatgg atggggttcc cttcacactc caccacgat ttgagggcaa gagctgcagc 420
cccctggcct tctctgcttt tggggacctg accatcaagt ctctggcgga cattgaggag 480

```

```

gagtataact acggcttcgt ggtggagaag accgcggctg cccgcctgcc cccagcgtc 540
tcatatgtccc tcacccttcc gcggaaagag ccccttact ccacctcccc gccagcctgg 600
ggccaccccc cctcactgca tcctgggaac cttcgccctg caaggcgttt gctatcttca 660
gccactgggc ggagctgcag ccctggagga gggggcgggt cgaggctgcg tggatgatgg 720
gtctccgccc ccacgccctg ccgggcaggg ctggagctgg acagaagcca gtgcctttaa 780
gtcattttgtg tcaaaaccct ctgggggtccg gaggtgtgtc ggggtgtcctc ctggcaataa 840
acactaccgg gttctcgcca aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa a
891

```

<210> 46

<211> 902

<212> DNA

<213> Homo sapiens

<220>

<223> clone 4S-6H8 - cDNA FLJ31039 fis, clone
HSYRA2000221

<400> 46

```

agagattctc taaatatgga attagattag gattctctgc tccacttaac acacattttt 60
aaattagtac tgatgattga gggatggaca atagcacacc aaaaaaaaaa agtttagtat 120
gaaaaattta aacctgttgg ttaagtcatt cccattaatg tcattttgct gagggtgact 180
tggtcctttt gaattgcttt ggtgtacggg tatgttctga tttttcatgc aagctcctct 240
gccattccac cgctctgagg agtaattgta gcacttcaca tgtgctgtgg ttgtgatcac 300
atggtgacat acatagcatg tgtgttccca gctgttgtgt gtttatgtga catttgatgc 360
caatacatat gtcttcaagg tatgcttgtt ccctcccagc togtggaata tcaaaaaaat 420
tcattgctgg aaaaattatt tcatagacaa aaatgttaat gttctcttgg ggacttagag 480
ttgaaaatat ttgtatagat ttggttctca agtccacaga atcgtatctg ctgtgggtctc 540
cctttgggtg tcactctgga gccatgtgta tggaagattc tgtcacaggc ggctgggatg 600
tgggcagatg ctgttagcct ccctctccac gtgggtgtcc atgcctgacg tgtcccctag 660
ttcaaggaag cgccatcttt agcatgaaaa caattgcgtt cccctaggaa atgaagaaaa 720
aatgagctga aatttcctta tacatttgaa tttgttcatt ttttaaagag acatttttgt 780
tgtctgcttt gtggtactta taaaatttgt tttccattga aattgccatt tataaatttg 840
cagatatgta ctaatttaga ttttttttaa gtgttcaata aaataaggat atatttactg 900
tg
902

```

<210> 47

<211> 566

<212> DNA

<213> Homo sapiens

<220>

<223> clone 4S-7F6 - cDNA FLJ13305 fis

<400> 47

```

aagactctct tagtgactct tccagatctg taccagaaaa gaactcctat caccctgtct 60
cattaatgac atcattttca gagcctgatt taggcagtc ttcctccttg tatgtgtcct 120

```

```

cctctgaaga ggagttaccc aacctagaaa aagagtatcc taggaaaaac agaatgatga 180
cctatgctaa ggagctcatc aacaatatgt ggacagactt ttgtgttgag gattatattc 240
gctgtaaaga tactggcttc catgcagctg aaaaaagaag gaagaaacga aaagaatggg 300
tgcccacaat tacagtaccg gagccttttc aaatgatgat aagagaactg aagaaaatct 360
tggtatggtg atgattttta ctctagtgcac agctgtgcaa gaaaaattaa atgaaatagt 420
agatcagata aaaactagaa gagaagaaga aaagaaacaa aaagaaaaag aagcagaaga 480
agctgaaaag caattattcc atggtactcc agttacaatt gagaatttct taaattggaa 540
agccaagttt gatgcagaac tcttgg                                     566

```

<210> 48

<211> 482

<212> DNA

<213> Homo sapiens

<220>

<223> clone 4S-7G6 - KIAA0776 protein (KIAA0776)

<400> 48

```

ttgatatcta ctgaaacata aatgataagg ttcttaaagg ttgaattaaa agtaatccat 60
gtttgtgtca aatgatcata gaaaataaat agaagagaca gtgaagcaag taaaaagaaa 120
agcattgttt taatttgttt gcattaattt ttttcatttg tcaaaatgct tcttttgttg 180
ccacagtaaa gaacagtttt tattgttttg taagtaaaat tacgtagctg attttgtatg 240
taaagattaa tttccataat aaaaattatt gtatgtttac tgtgatctta atgggcaggg 300
ttaagaaagt tatttaaaat aaagttacct attctactaa attttatagt actttgaagc 360
ttctattaat taacacaaag attaattggg gcataatatt tatatatata ctttttgaat 420
tctcattttg aacattatta aaggatttta ttttctttac acaaaaaaaaa aaaaaaaaaa 480
aa                                                                 482

```

<210> 49

<211> 274

<212> DNA

<213> Homo sapiens

<220>

<223> clone 4S-9F4 - similar to hypothetical protein
FLJ10751

<400> 49

```

agatgagggg taggtgtgcc cagccctcca gacccggcct ttctggttaa cccctgcatg 60
ccaagctgcc tgctgcccc ggtcctcacc tcaggccttt gaaggggcag cttctggaag 120
ttgttttctc ctctgcttgg agagtttgcc cttgtctgtc ttggaaagtg tgggcagcca 180
cagatgcccc caaatcagag ctacacagtga gtgagcccct aagcttcagt ctgcaataaa 240
gaatgcattg gtttcatcaa aaaaaaaaaa aaaa                                     274

```

<210> 50

<211> 1021

<212> DNA

<213> Homo sapiens

<220>

<223> clone 4S-9G10 - hypothetical protein BC013073
(LOC92703)

<400> 50

```

tgagagaact gtctccagaa aaaagaaaag caagagacac aaagaagaac tggacggggc 60
tggaggagaa gagtatccca tggatatattg gctattgctg gcctcctata tccgtcctga 120
ggacattgtg aatTTTTccc tgatttgtaa gaatgcctgg actgtcactt gcaactgctgc 180
cttttgacc aggttgtagc gaaggcacta cacgctggat gcttcctgc ctttgctgtc 240
gcgaccagag tcaatggaga agctgcgctg tctccgggct tgtgtgatcc gatctctgta 300
ccatatgtat gagccatttg ctgctcgaat ctccaagaat ccagccattc cagaaagcac 360
ccccagcaca ttaaagaatt ccaaagtctt acttttctgg tgcagaaaga ttgttgggaa 420
cagacaggaa ccaatgtggg aattcaactt caagttcaaa aaacagtccc ctaggttaaa 480
gagcaagtgt acaggaggat tgcagcctcc cggtcagtac gaagatgttc ataccaatcc 540
agaccaggac tgctgcctac tgcaggcac caccctcaat ttcattctta ttccgattgt 600
catgggaatg atatttactc tgtttactat caatgtgagc acggacatgc ggcatcatcg 660
agtgagactg gtgttccaag attcccctgt ccatgggtgt cggaaactgc gcagtgaaca 720
gggtgtgcaa gtcatcctgg acccagtgc cagcgttcgg ctctttgact ggtggcatcc 780
tcagtaccca ttctccctga gagcgtagtt actgcttccc atcccttggg ggcagcctcg 840
agtgtagtc attagtaatc agattccagt ttggacaggg tggctggatt gtatatctcg 900
ttagtaatgt acatgctctt caggttctag ggctcctgtt aggggaggga gaaatgttga 960
atcaagaggg aaaacaacta ctatgattta taaacatatt ttaatgtaaa aatttgcatt 1020
t

```

1021

<210> 51

<211> 1343

<212> DNA

<213> Homo sapiens

<220>

<223> clone 5S-21/57 - hypothetical protein FLJ10081

<400> 51

```

agatgattga cagtgaactc actatgcagg gctgttggt ccaacctgag ccctataggt 60
ggcagtcctt ggagaagtgg tcacagaaga tggagctctg atccctgct tacctcttca 120
caacacttgt gtgcaaagat agtttttagat ttggtttaga agctatcctc cagaacaggc 180
tcccatactt agaatgtttc tagttaagggt aataaattag gcaaccaag tgtgactcca 240
ctcaagtgtc cttttctgta ggcaggaagg gccacaaca tggcttaaaa tgtagtccat 300
ggttctggcc cacagtacag tgtgtatcta taccagggtc cctgtgttca atctgggagc 360
cttcctggcc agtctgagtg gcagccagaa gggagctcat agtgtctagg agtctcargc 420
aaggtaggtc aggttactgt gggcaggggg gatgtgtgtg ataggagagg gtaccctaaa 480
ccccatacct tccctccctg acctgaaaaa gctgatcttc aacagggtt cacacagaat 540
taggctgtgt ttttgcatta gctggtagggt gactttctca aaattcttaa attcagaaag 600

```

tatttanta aa acttgaggaa ggtatgaaat ctggaggagg catccaggac ccagggggtt 660
 gatagcttta caggtaggat catacccacc caaaagagca gtggacaata agactatttg 720
 agctatatga agcttttagg aatcatttag gacagacaga gcccttaamc aaccattca 780
 tgacttaagt tgttggttc artgtwtkcc tggggacaaa gaaaaactaa caagcccgac 840
 ctgcctttat gataaattct agtgtgctwa caagggatga cttcctgagg tgtgatctgt 900
 ccaccttgaa gaactccaca actgaagaag gggagctgtg agaacgtgga ttgttctaca 960
 actgacagag ggtaacagag gaagtggctg aggcctagag tcacgttttc cagttccctt 1020
 cgcaaactat atttcttgga acgcgaaagg aagctttacc tatttcataag aagacctgga 1080
 atccataacc tcagaaggca atattattga tagaaaatgt ggaaggatca ggaagttctt 1140
 agattcttgg atgacanatg catgttgatg ccctatggag atgtccttgt gttttgaggt 1200
 cactgaggta ggaagacctg tctactcttg gtttcaccac tagaacagtc ttgggctgga 1260
 tgggttatag agctgagcgg ctgtgatggt tctgttttta cattaacaaa aacaattaaa 1320
 aacaccaaaa acaaaaaaaaa aaa 1343

<210> 52

<211> 2073

<212> DNA

<213> Homo sapiens

<220>

<223> clone 8C10 - hypothetical protein FLJ23018 (
 FLJ23018)

<400> 52

aagaaaattc gagggaaaga agtttacatg actatggctt acggcaaggg agaccccctc 60
 cccccacca ggctgcagca cagtatgcat tatgggcacg atcctccaat gcactactca 120
 cagacagctg gcaatgttat gtctaataaa cattttcatc ctcagcatcc atctccgaga 180
 caaggctcgg gatatgggat gcccaggaat tcatctcggg ttataaacag gcacaacatg 240
 ccggggcccta aagttgattt ttaccaggc ccaggtaaaa ggtgctgcca gagctatgat 300
 aacttctctt atagatctcg ttcattttaga cgtagtcacc gccagatgag ttgtgtgaat 360
 aaggagtccc agtatggatt taccaggc aatggacaga tgcccagggg cttggaagaa 420
 actattactt tttatgaagt tgaagaagg gatgagactg cttatgcaac ttttcctaata 480
 catgaaggct cctctacaat ggttcctgct acttcaggat actgtgttg aaggcgggga 540
 catagctcag gcaaacagac tttgaattta gaggagggca atggccagag tgaaaatggg 600
 cgatatcatg aagaatatct ttatcgtgca gagccagact atgaaacttc aggtgtttat 660
 agcacaactg catctacagc aaacttgtct cttcaggaca gaaagtcagc ttctatgtct 720
 cctcaggaca cagttacctc atacaactac cccagaaga tgatgggaaa tattgcagca 780
 gttgcagctt cctgtgcaa taatgttcca gctccagctc tatctaaccg tgcagcggct 840
 aatcaagcta ttagtaccac ttcagtttcc tcacagaatg ctatacagcc tctctttgta 900
 tctccaccta cacacggcag gccagatata aaagttttgc agtactattt caatctagga 960
 ttgcagtgtc attaccacag ctactggcac tccatggctc atgtgccaca gatgcagcag 1020
 cagcttcatg tagagaatta tccagtctat actgagccac ctctggtaga tcaaaccgtt 1080
 cctcaatgct acagtgaggt gaggagagaa gatggcatac aggcggaagc atcagcaaat 1140
 gatacttttc cgaatgctga ttcttcatct gtccctcatg gagcagctca ttatccagta 1200
 atgtcagatc cctatgggca gccaccttg ccaggttttg actcctgcct tccggttggt 1260
 ccagattatt cctgtgttcc cccctggcat ccagttggta cagcatatgg tggttcttct 1320
 caaattcatg gtgctataaa tccctgggcca attggctgta ttgctccatc tccccagct 1380


```

tctcattatg tacctcaggg tatgtaagat ccagcagtat gaagtattct tgcactgcc 1440
ttttcttgct gtttttgttt ttaaaaagta ttttatgtta gtggttaaat gatttaggtg 1500
attagtgttt actattgtat ttgtctttta aattatttta tcttttgatt taaaatagta 1560
ctttaaaatt aaggggtatt attttgggct gtgactaagg aaattgagat ggatgtacaa 1620
ctagcccat attgagcata cttcattgta ttcagctgtt ttcctgtcag ccatttgtca 1680
gctttatatt agctgatgtt accaattgat aaaatgaata taaagtattt cattggttca 1740
aaaatcacac atcatattaa accatgcaga attggagtaa cttccacttt tttctagaaa 1800
gtaaaaccaa gagcctttgc ttctggataa ctacttaaat attaaattaa agagctcttc 1860
acgtttcttg agaattatct gaagccagtt gcattctgtg atatcagttt tgaaggcaca 1920
tggttctctg ctttagattt atcccatatg ctattgttta atactggatg tatgtaagtg 1980
ttttactgca ctgtattgaa ttggtgtctt ttgcacagtt agcagtaaat aaaaattagc 2040
atttaaaatt gcaaaaaaaaa aaaaaaaaaa aaa 2073

```

<210> 53

<211> 804

<212> DNA

<213> Homo sapiens

<220>

<223> clone 5D-15/114 - cDNA DKFZp4340159

<400> 53

```

tatgattgaa atgcattcat tcatcacgca taggcacaat cacaacttga tgatgcttgg 60
gaaagaatca acagttaaaa cttcatgaag ttctaattgtc tgtgttccaa aacacatcac 120
attattaggt ttaggggaca tacgtagggtg tgctccctgg ggtggggagt tttctagtta 180
ctagaccatc tcccattttt agcacttggc agcctcatga tccttttata aataggagat 240
taacaggaga gcagcaatac gattttgcca atggaataac agatttgccg gcattcactg 300
aaagaggggca satattgggt ccttgtgact tcaactgact cttccgaatt gtatgaattt 360
atcaatgtat tagataaacc cagtttcaga ataataaaga aaaaatatta gaccaaataa 420
tgtggcta atgtgggtatg atttctagcc cgtgggttta aaactgtatc ctaaagagtc 480
attttaaaat aatatwaata tttaaaaatg taactgctat ctttatgttc tgaaataagt 540
taaaacattt taaaatatga atactgtagt ttaaaagaaa gaaatggtgg gaaggaaaag 600
tagagaaaga aatgcccaatt ccagtccaaa gctttgtttg ccaagttttc ttasaatgaa 660
ttttaccaat gtatgggttc ttgttaacag aatgtgtaac agaaatactg aaagactttt 720
gcctaaagtg gcattattga ctgctggtgt gatgctactg taatgcgata aattattaaa 780
ttgttgcaaa gtgaaaaaaaa aaaa 804

```

<210> 54

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer used
for generation of a ferritin PCR fragment

<400> 54

ctacgagcgt ctctgaaga tgc

23

<210> 55

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer used
for generation of a ferritin PCR fragment

<400> 55

cgcggtacca agtcgctggg ctcaagaaggc tc

32

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
5 June 2003 (05.06.2003)

PCT

(10) International Publication Number
WO 2003/045988 A3

(51) International Patent Classification⁷: **C07K 14/39**,
C12N 15/10

(21) International Application Number:
PCT/EP2002/013549

(22) International Filing Date:
28 November 2002 (28.11.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
01204600.9 29 November 2001 (29.11.2001) EP

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*):
**VLAAMS INTERUNIVERSITAIR INSTITUUT
VOOR BIOTECHNOLOGIE VZW** [BE/BE]; Rijviss-
chestraat 120, B-9052 Zwijnaarde (BE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **CONTRERAS,**
Roland, Henry [BE/BE]; Molenstraat 53, B-9820 Merel-
beke (BE). **CHEN, Cuiying** [BE/BE]; Lijsterstraat 10,
B-9860 Balegem (BE).

Published:

— with international search report

(88) Date of publication of the international search report:
4 March 2004

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 2003/045988 A3

(54) Title: METHOD TO ISOLATE GENES INVOLVED IN AGING

(57) Abstract: The present invention relates to a method to isolate genes involved in aging and/or oxidative stress, by mutation or transformation of a yeast cell, subsequent screening of the mutant or transformed cells that are affected in aging and isolation of the affected gene or genes, and the use of these genes to modulate aging and aging-associated diseases in a eukaryotic cell and/or organism.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 02/13549

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07K14/39 C12N15/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, EMBL, BIOSIS, MEDLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 01 21786 A (KEIO UNIVERSITY ;NISHIMOTO IKUO (JP)) 29 March 2001 (2001-03-29) see example 2, p. 37, example 4, p. 40, example 6, p. 41 ----- -/--	8,13

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

12 March 2003

Date of mailing of the international search report

24. 06. 03

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Paresce, D

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/13549

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	MAX INGMAN, HENRIK KAESMANN, SVANTE PÄÄBO & ULF GYLLENSTEN: "Mitochondrial genome variation and the origin of modern humans" NATURE, vol. 408, no. 6813, 7 December 2000 (2000-12-07), pages 708-713, XP002190961 see abstract	8,13
X	-& DATABASE EMBL 'Online! 14 March 2001 (2001-03-14) "Homo sapiens mitochondrion, complete genome" Database accession no. AF346979 XP002234371 abstract	8,13
Y	--- HASHIMOTO Y., ET AL.: "A rescue factor abolishing neuronal cell death by a wide spectrum of familial Alzheimer's disease genes and A beta" PROC. NATL. ACAD. SCI. U.S.A., vol. 98, no. 11, 22 May 2001 (2001-05-22), pages 6336-6341, XP002190962 page 6336	8,13
X	-& DATABASE EMBL 'Online! "Homo sapiens Humanin (HN1) mRNA" Database accession no. AY029066 XP002234372 abstract	8,13
Y	--- US 2001/026930 A1 (AUSTRIACO NICANOR ET AL) 4 October 2001 (2001-10-04) see p. 9, right column, claims 1, 6	1-6,8,9, 13
Y	--- US 5 919 618 A (KENNEDY BRIAN ET AL) 6 July 1999 (1999-07-06) see columns 4-5, 16-17 and claims	1-6,8,9, 13
Y	--- WO 95 05459 A (MASSACHUSETTS INST TECHNOLOGY ;GUARENTE LEONARD P (US); AUSTRIACO) 23 February 1995 (1995-02-23) cited in the application see p. 28-33, claims 2, 10, 18, 26-33 --- -/--	1-6,8,9, 13

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/13549

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>GRZELAK A ET AL: "Decreased antioxidant defense during replicative aging of the yeast <i>Saccharomyces cerevisiae</i> studied using the 'baby machine' method"</p> <p>FEBS LETTERS, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 492, no. 1-2, 9 March 2001 (2001-03-09), pages 123-126, XP004257348 ISSN: 0014-5793 see materials and methods and p. 124, left column</p> <p>-----</p>	1-6,8,9, 13

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 02/13549

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 9-10, 13-15 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: 7, 11-12
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-6 completely, 8-9, 13 partially

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

Invention 1: claims 1-6 completely, claims 8-9,
13 partially

Claims 1-6 completely, claims 8-9, 13 partially are directed to a method to screen genes involved in aging, as well as genes isolated by said method, in particular a gene comprising SEQ ID NO: 1, the polypeptide expressed from said gene, and uses thereof.

Invention 2: claims 8-9, 13 partially

Claims 8-9, 13 partially are directed to a gene comprising SEQ ID NO: 3, the polypeptide expressed from said gene, and uses thereof.

Invention 3-45: claims 8-10, 13-15 partially

Inventions 3-45 are directed to genes or gene fragments involved in aging or oxidative stress. Each claimed sequence is considered a separate invention, the inventions are numbered according to the list of sequences given in claim 8, namely, invention 3 is directed to the gene comprising SEQ ID NO: 5, invention 4 is directed to the gene comprising SEQ ID NO: 7, etc. Note: claim 10 is searched partially for invention no: 7 and 10, claim 14 is searched for invention no: 7 and claim 15 is searched for invention no: 10. The order of inventions follows the order of sequences given for claim 8. The inventions are also directed to the polypeptides expressed from said genes, and uses thereof.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 7, 11-12

Present claims 7, 11-12 and claim 9 partially relate to an extremely large number of possible genes or gene fragments. The claimed genes are characterized only by the fact that they may be isolated by a certain method. However, support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds/products claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those claims which appear to be supported and disclosed, namely those that refer to specific sequences or methods (claims 1-6, 8, 10, 13-15 and claim 9 partially).

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 02/13549

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0121786	A	29-03-2001	AU 7313800 A	24-04-2001
			AU 7313900 A	24-04-2001
			CA 2385444 A1	29-03-2001
			EP 1221480 A1	10-07-2002
			WO 0121786 A1	29-03-2001
			WO 0121787 A1	29-03-2001
<hr/>				
US 2001026930	A1	04-10-2001	US 6218512 B1	17-04-2001
			US 5919618 A	06-07-1999
			US 5874210 A	23-02-1999
			WO 9505459 A1	23-02-1995
<hr/>				
US 5919618	A	06-07-1999	US 6218512 B1	17-04-2001
			US 2001026930 A1	04-10-2001
			US 5874210 A	23-02-1999
			WO 9505459 A1	23-02-1995
<hr/>				
WO 9505459	A	23-02-1995	WO 9505459 A1	23-02-1995
			US 6218512 B1	17-04-2001
			US 5919618 A	06-07-1999
			US 2001026930 A1	04-10-2001
			US 5874210 A	23-02-1999
<hr/>				

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☒ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)